

Combined effect of bicarbonate and water in photosynthetic oxygen evolution and carbon neutrality

Yanyou Wu¹ 

Received: 21 October 2022 / Revised: 21 October 2022 / Accepted: 1 November 2022 / Published online: 18 November 2022
© The Author(s), under exclusive licence to Science Press and Institute of Geochemistry, CAS and Springer-Verlag GmbH Germany, part of Springer Nature 2022

Abstract Carbon neutrality is widely concerned and highly valued by many countries. Biosphere has always maintained the balance between oxidized organic substances and assimilated organic matter, resulting in net-zero carbon dioxide (CO_2) emissions and maintaining its own carbon neutrality. Nature has set a good example for human beings to coordinate oxygen (O_2) balance and CO_2 balance, and achieve carbon neutrality. How does photosynthetic oxygen evolution initiate carbon and water neutrality? My synthesis shows that photosystem II functions as carbonic anhydrase to catalyze the reaction of CO_2 hydration under physiological conditions, and CO_2 hydration coupled with chemical equilibrium, $\text{H}^++\text{HCO}_3^- \rightarrow 1/2 \text{O}_2 + 2\text{e}^- + 2\text{H}^+ + \text{CO}_2$, occurs in a photosystem II core-complex. Meanwhile, I focused on the revisiting of four classical heavy oxygen (O^{18}) labeling experiments and found that bicarbonate can promote photosynthetic oxygen evolution, and that photosynthetic oxygen evolution can alternately come from bicarbonate and water, not only water. Bicarbonate photolysis and water photolysis account for half of the photosynthetic oxygen evolution respectively, which can well explain the bicarbonate effect, Dole effect and plants' environmental adaptability. Photosynthetic oxygen evolution initiated the journey of water metabolism and carbon metabolism in nature, which led to the coupling as 1:1 (mol/mol) stoichiometric relationship between the reduction of CO_2 and oxidation of organic

carbon, coordinated the evolution of the atmosphere, hydrosphere, lithosphere and biosphere, and realized "carbon neutrality" in the whole Earth system.

Keywords Bicarbonate photolysis · Bicarbonate effect · Carbonic anhydrase · Dole effect · Water photolysis · Stoichiometric relationship · Carbon metabolism

1 Introduction

Carbon neutrality is a widely discussed topic worldwide. Photosynthesis, which converts light energy into chemical energy and inorganic carbon into organic carbon to releases oxygen, is the most important biochemical reaction on Earth and the core link of carbon neutrality. The evolution of photosynthetic oxygen plays a crucial role in photosynthesis. Understanding the photosynthetic oxygen evolution can provide a solution to carbon neutrality in humans.

In 1937, Hill confirmed that in vitro chloroplasts could reduce trivalent iron compounds to divalent iron compounds and release oxygen under light (Hill 1937). Although there have been some disputes, many heavy oxygen (O^{18}) isotope labeling experiments have led to a general consensus that the released oxygen comes only from water, which seems to be "conclusive" isotopes evidence (Ruben et al. 1941; Warburg and Krippahl 1958; Stemler and Radmer 1975; Radmer and Ollinger 1980; Stemler 1980, 2002; Hillier et al. 2006). However, can these experiments really prove that the released oxygen during photosynthesis comes solely from water, is excluded from bicarbonate? Let us revisit these heavy oxygen isotope labeling experiments.

 Yanyou Wu
wuyanyou@mail.gyig.ac.cn

¹ State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, 550081 Guiyang, People's Republic of China

2 Reappearance of scenes

In 1941, Ruben et al. reported the results of a heavy oxygen (O^{18}) labeling experiment of photosynthesis. In this study, *Chlorella* cells produced heavy oxygen under light conditions when the algae were suspended in heavy oxygen water (0.85% O^{18}) containing ordinary potassium bicarbonate and carbonate. However, *Chlorella* cells were suspended in ordinary water and heavy oxygen potassium bicarbonate and carbonate, and the algae evolved unlabeled oxygen in the presence of light. Therefore, Ruben et al. concluded that oxygen originated solely from water according to isotope ratios identical to those of water (Ruben et al. 1941). Many other scientists have reached a similar conclusion (Govindjee et al. 2006).

This conclusion was questioned shortly afterwards. In 1958, Warburg and Krippahl first discovered that bicarbonate can stimulate *Chlorella* to release oxygen under the action of artificial reductants, which was later called the bicarbonate effect, and bicarbonate or CO_2 was necessary for the Hill reaction. They believed that the molecular oxygen of the Hill reactions is evolved not from the splitting of water, but from the splitting of the hydrocarbon (Warburg and Krippahl 1958).

The bicarbonate effect in the Hill reaction has been widely confirmed. However, scientists have proven that oxygen still comes only from water photolysis in the Hill reaction under the stimulation of bicarbonate using $HC^{18}O_3^-$ as a tracer. In 1975, Stemler and Radmer conducted an $HC^{18}O_3^-$ labeling experiment using disrupted chloroplasts depleted of bicarbonate/ CO_2 and lacking the enzyme carbonic anhydrase (CA) to determine whether bicarbonate acts as the immediate source of O_2 . They found that HCO_3^- -depleted broken chloroplasts evolved only small amounts of ordinary O_2 without the addition of $HC^{18}O_3^-$ under illumination. However, the HCO_3^- -depleted broken chloroplasts with the addition of $HC^{18}O_3^-$ under illumination released oxygen that was almost entirely unlabeled and less than 0.1% was doubly labeled. Therefore, they concluded that HCO_3^- was not the immediate source of photosynthetically evolved oxygen, as the administration of bicarbonate produced a fivefold stimulation of oxygen evolution in the chloroplast Hill reaction (Stemler and Radmer 1975).

Single short saturating light flash experiment demonstrated that the evolution of oxygen was accompanied by flash-induced period-four oscillations. The dark-adapted chloroplasts were subjected to the first and second flashes, and little or no O_2 was produced. After the third flash, O_2 was released at most. After the fourth flash, the evolution of O_2 was the second highest, followed by a gradual decrease to a constant value (Allen and Franck 1955; Kok et al. 1970; Joliot 2003). The above flash-induced photosynthetic O_2 has also been experimentally proven that it came only from

water photolysis. In 1980, Radmer and Ollinger determined the evolution of O_2 elicited by short saturating light flashes in HCO_3^- -depleted chloroplasts in the dark reactivated with $HC^{18}O_3^-$. They found that “all” the O_2 evolved was unlabeled. Therefore, they supported that the O_2 evolved from neither bicarbonate nor CO_2 (Radmer and Ollinger 1980).

By the 21st century, many scientists still insisted that water was the sole substrate for O_2 production by photosystem II in contemporary oxygenic organisms. For example, in 2006, Hillier et al. measured the ^{18}O enrichment of the O_2 produced in 50 mM bicarbonate enriched to 96% ^{18}O and 50 mM unlabeled bicarbonate both with 10 μM ethoxzolamide (EZ) (a CA inhibitor) for spinach PS II membrane fragments illuminated by 16 saturating light flashes. They found that there was a small (~ 5% relative) increase in the ^{18}O enrichment of the O_2 evolved in the presence of labeled bicarbonate compared with that measured in 50 mM unlabeled bicarbonate. Almost all photosynthetic oxygen that evolves in the presence of labeled or unlabeled bicarbonate is unlabeled oxygen. Accordingly, they concluded that bicarbonate is not a physiologically significant substrate, and the bulk of O_2 produced comes from water; this is because exceedingly low levels of bicarbonate oxidation may occur in contemporary photosystem II (Hillier et al. 2006). The above-mentioned heavy oxygen labeling assessments give the impression that O_2 evolving only from water photolysis is irrefutable (Fig. 1).

3 Challenge


3.1 Challenge 1 bicarbonate significantly influences photosynthetic oxygen evolution

Bicarbonate effect in the Hill reaction is the most effective argument against the fact that O_2 evolved only from water photolysis. Oxygen evolution can increase several times or more than ten times under the stimulation of bicarbonate; however, oxygen evolution of *Chlorella* illuminated completely cannot be begin without inorganic carbon (Warburg and Krippahl 1958; Warburg et al. 1965), which demonstrates that the premise of oxygen evolution should be in the presence of inorganic carbon. The role of bicarbonate played in photosynthetic oxygen evolution is significantly greater than that of water.

3.2 Challenge 2 existence of Dole effect is strongly conflict with that the oxygen comes only from water photolysis

If the oxygen released by plants during photosynthesis only comes from water, the oxygen isotope composition of oxygen in the atmosphere should be very close to that of seawater. However, atmospheric O_2 is nearly 24‰ more

Fig. 1 Schematic diagram of photosynthetic oxygen evolution that is generally accepted and written into textbooks. The oxygen evolved comes only from water photolysis; that is, 2 moles of water yield 1 mol of oxygen. Photosynthetic oxygen evolution is not associated with CO₂ assimilation



enriched in ¹⁸O than seawater, and this enrichment is known as the Dole effect (Metzner 1975). Even if scientists take all factors (including physical, chemical and biological factors) into account, the global ¹⁸O enrichment in atmospheric O₂ with respect to ocean water (Dole effect) would still be 21‰–24‰ (Bender et al. 1994; Hoffmann et al. 2004; Mader et al. 2017). This indicates that the sources of oxygen in the atmosphere is not only water, but also another source almost equal to water source.

3.3 Challenge 3 thermodynamically, bicarbonate photolysis is more feasible than water photolysis in photosystem II

The standard free energy variance in the chemical equilibria (H⁺+HCO₃⁻ → 1/2 O₂ + 2e⁻+2 H⁺+CO₂) was 24.8 kcal/mol, and that in the chemical equilibria (H₂O → 1/2 O₂ + 2e⁻+2 H⁺) was 37.3 kcal/mol (Dismukes et al. 2001). The free energy required for O₂ evolution using bicarbonate as the substrate is significantly lower than that required using water as the substrate during photosynthesis of organisms. This result demonstrates that bicarbonate photolysis is more likely to occur during photosynthetic O₂ evolution than water photolysis.

3.4 Challenge 4 thylakoid carbonic anhydrase (CA) and photosystem II with CA activity

3.4.1 Many types of carbonic anhydrases in chloroplasts

Carbonic anhydrase (CA, EC 4.2.1.1) has many physiological functions and is ubiquitously found in most cells

from all kingdoms of life. CA catalyzes the reversible conversion of bicarbonate to CO₂, which is 10⁷ times the rate of non-enzymatic reaction and one of the fastest enzymatic reactions. Many types of CA have been found in chloroplasts. CA has also been found tightly bound to thylakoid membranes, and is called thylakoid CA (tCA) (Pronina et al. 2002; Moskvin et al. 2004; Rudenko et al. 2006, 2007).

Thylakoid CA (tCA) shows an association between enzymatic activity and photosystem II (Khristin et al. 2004; Ignatova et al. 2011; Rudenko et al. 2015). On the one hand, the CA inhibitors, acetazolamide and trifluoromethanesulfonamide, suppress not only CO₂ hydration but also electron transfer and oxygen evolution in photosystem II. In addition, this inhibition was completely reversed by the addition of bicarbonate (Shitov et al. 2011, 2018). Meanwhile, bicarbonate functions not only as a mobile proton acceptor, but also as a light-driven producer of both O₂ and CO₂ (with a constant stoichiometric relationship) (Koroidov et al. 2014). This indicates that bicarbonate is involved in electron transfer as proton acceptor, electron donor, oxygen and CO₂ evolution in the photosystem II. On the other hand, tCA activity is dependent upon the surrounding redox potential and is similar to that of photosystem II (Moubarak-Milad and Stemler 1994). Similarly, photosystem II inhibitors, hydroxylamine and 3-(3,4-dichlorophenyl)-1,1'-dimethylurea (DCMU), inhibit the activity of tCA (Rudenko et al. 2015). This indicates that tCA catalyzes a redox reaction similar to the photosystem II.

3.4.2 Photosystem II with CA function differently from that of typical CA

To date, the CA located in the photosystem II core-complex cannot be completely isolated, thus, it is called intrinsic CA. The intrinsic CA in the photosystem II core-complex undergoes only CO_2 hydration under physiological conditions (Lu and Stemler 2007). Some inhibitors exhibit good inhibitory effects on the activity of CA, but have little effect on oxygen evolution in the photosystem II (Rodionova et al. 2017). This may indicate that CA is not a separable part of the photosystem II core-complex. Photosystem II likely functions both as oxygen evolution and CA, however, it is not as a typical CA.

3.4.3 The activity of oxygen evolution in the tCA/photosystem II is dependent on Mn

The Mn clusters (Mn_4CaO_5) of photosystem II, as oxo bridges linking the Mn atoms, may be catalyzed for dioxygen formation to yield oxygen (Umena et al. 2011). The oxygen evolution activity of photosystem II core-complex depends on the presence of Mn. The activity of oxygen evolution was inhibited, but the activity of CO_2 hydration of photosystem II core-complex was not suppressed after removing Mn clusters (Dai et al. 2001). Moreover, CA Mn-substituted the active-site zinc (CA[Mn]) functions as peroxidase and produces O_2 in the presence of bicarbonate (Okrasav and Kazlauskas 2006). This implied that, like CA[Mn], photosystem II core-complex causes bicarbonate oxidation to produce oxygen catalyzed by Mn. The above interaction between the photosystem II and bicarbonate suggests that CO_2 hydration is coupled with the chemical equilibria, $\text{H}^+ + \text{HCO}_3^- \rightarrow 1/2 \text{O}_2 + 2\text{e}^- + 2\text{H}^+ + \text{CO}_2$, and occurs in a photosystem II core-complex.

3.5 Challenge 5 sulphid photolysis in anaerobic photosynthetic bacteria during photosynthesis

Some anaerobic photosynthetic bacteria, such as *Oscillatoria limnetica*, split hydrogen sulfide rather than water, and release sulfur rather than oxygen. Thus indicates that photosynthetic organisms can split substances other than water during photosynthesis under light (Cohen et al. 1975). Thus, water is not the sole substrate used for photolysis during photosynthesis.

4 Revisiting some heavy oxygen labeling experiments

4.1 Revisiting the heavy oxygen (O^{18}) labeling experiment from Ruben et al. (1941)

This experiment was performed using active *Chlorella* cells. Algae were suspended in ordinary or heavy oxygen water and heavy oxygen or ordinary potassium bicarbonate and carbonate. They found that the $\text{O}^{18}/\text{O}^{16}$ ratio of oxygen evolved during photosynthesis by *Chlorella* is identical to that of water. Therefore, they concluded that the oxygen evolved by *Chlorella* originated only from water. The premise of this conclusion drawn is that the oxygen in group such as OH, C=O, COOH exchanges very slowly with water under the experimental conditions (Ruben et al. 1941).

However, the premise does not occur in the experiment. *Chlorella* has many types of CAs (Hogetsu and Miyachi 1979), and photosystem II in algae inevitably function as CAs. CA rapidly catalyzes the chemical equilibria $\text{H}^+ + \text{HCO}_3^- \leftrightarrow \text{H}_2\text{O} + \text{CO}_2$, thereby speeding up the exchange of heavy oxygen in $\text{HC}^{18}\text{O}_3^-$ with ordinary oxygen in water, or ordinary oxygen in bicarbonate with heavy oxygen in H_2^{18}O . The maximal concentration of bicarbonate used in the experiment was lower than 0.2 M, which was less than 0.4% of the water concentration (55.6 M) in the system. The oxygen in bicarbonate would exchange with almost all oxygen in water. Finally, irrespective of whether water photolysis or bicarbonate photolysis occurs in the experimental system, the $\text{O}^{18}/\text{O}^{16}$ ratio of oxygen evolved in photosynthesis by *Chlorella* would always be found identical to that of the water (Fig. 2a).

4.2 Revisiting the heavy oxygen (O^{18}) labeling experiment from Stemler and Radmer (1975)

This experiment was performed using HCO_3^- -depleted maize chloroplast fragments. Stemler and Radmer found that HCO_3^- -depleted broken chloroplasts released oxygen stimulated by $\text{HC}^{18}\text{O}_3^-$ under illumination, which was almost entirely unlabeled and less than 0.1% doubly labeled. Similarly, they concluded that the photosynthetically evolved oxygen stimulated by bicarbonate also comes from water rather than inorganic carbon. This conclusion is reached on the premise of excluding the influence of exogenous inorganic carbon and effectively controlling the isotope exchange between H_2O and CO_2 (Stemler and Radmer 1975).

However, although researchers removed nearly all CAs and HCO_3^- from broken chloroplast fragments, this experiment did not isolate CO_2 from the air, or effectively




Fig. 2 Bicarbonate photolysis was performed to explain two classic heavy oxygen labeling experiments under continuous light. **(a)** In the experiment from Ruben et al. (Ruben et al. 1941), carbonic anhydrases (CAs) in *Chlorella* rapidly catalyzes CO_2 hydration and bicarbonate dehydration, thereby speeding up the exchange of heavy oxygen in $\text{HC}^{18}\text{O}_3^-$ with ordinary oxygen in water, and also the exchange of ordinary oxygen in bicarbonate with heavy oxygen in H_2^{18}O . The concentration of water is much greater than that of inorganic carbon, and the oxygen in bicarbonate exchanges with almost all the oxygen in water. **(b)** In the experiment from Stemler and Radmer (Stemler and Radmer 1975), the action of photosystem II (like CA) causes the exchange of the oxygen in OH , $\text{C}=\text{O}$ and COOH . Since the amount of unlabeled CO_2 in the air and the amount of water as a solvent are respectively much greater than the amount of unlabeled CO_2 and labeled H_2O converted from labeled bicarbonate, almost all bicarbonates used for photolysis are unlabeled

control the group exchange of oxygen among OH , $\text{C}=\text{O}$ and COOH in a bicarbonate-stimulated Hill reaction system. When $\text{NaHC}^{18}\text{O}_3$ was added into the Hill reaction system of HCO_3^- -depleted broken chloroplasts during the

dark period, they obtained the maximum signal of C^{18}O_2 and recorded a certain amount of ordinary signal of CO_2 within a few seconds. This indicates that the conversion of bicarbonate to CO_2 and H_2O and the exchange of labeled

CO_2 and unlabeled CO_2 were performed in the Hill reaction system. Because CO_2 is a liposoluble substance, it can easily enter the thylakoid membrane. However, the water-soluble bicarbonate remaining in the Hill reaction system could not pass through the thylakoid membrane. After the onset of illumination, the photosystem II in the thylakoid membrane undergoes extremely rapid CO_2 hydration to form bicarbonate, which then quickly transformed into H_2O and CO_2 under physiological conditions (Lu and Stemler 2007). The action of photosystem II (like CA) lasts for a long time (more than seconds) (Delosme and Joliot 2002), which almost leads to a full exchange of the oxygen in CO_2 with the oxygen in H_2O , resulting in minimal heavy oxygen (O^{18}) labeling $\text{HC}^{18}\text{O}_3^-$. Therefore, if the photolysis of bicarbonate occurs in HCO_3^- -depleted maize chloroplast fragments, the bicarbonate as the substrate is barely labeled, and the oxygen evolved would also be almost entirely unlabeled (Fig. 2b).

4.3 Revisiting the heavy oxygen (O^{18}) labeling experiment from Radmer and Ollinger (1980)

The experiment was performed using HCO_3^- -depleted chloroplasts under the action of short saturating light flashes. Radmer and Ollinger found that oxygen evolved by the HCO_3^- -depleted chloroplasts exhibited a 7- to 8-fold decrease in the rate of the Hill reaction via the untreated chloroplasts under continuous light. Meanwhile, the species and magnitude of O_2 evolved by HCO_3^- -depleted chloroplasts were similar to those evolved by depleted chloroplasts reactivated with unlabeled or labeled HCO_3^- under the action of short saturating light flashes. In fact, the experiment indicated that the oxygen evolved by HCO_3^- -depleted chloroplasts under short saturating light flashes was one order of magnitude less than that under continuous illumination, while being independent of exogenous HCO_3^- used for activation (Radmer and Ollinger 1980). However, the existence of bicarbonate photolysis by photosystem II was not ruled out.

Many studies have shown that a considerable amount of bicarbonate is bound to the vicinity of the oxygen-evolving center of the photosystem II, either (both) in the acceptor or (and) in the donor side of photosystem II, and it seems to flow freely (Van Rensen and Xu 1999; Stemler 2002; Shevela et al. 2012; Koroidov et al. 2014; Tikhonov et al. 2018). However, bound bicarbonate in the photosystem II is not necessary for photosynthetic O_2 evolution (Van Rensen and Vermaas 1981), and many binding sites for bicarbonate do not always exist, depending on the pH value (Bowden et al. 1991). The architecture of the photosynthetic oxygen-evolving center does or does not display bi(tri)dentate ligand-bound bicarbonate in different observations (Ferreira 2004; Umena et al. 2011). These results


indicate that bicarbonate bound in photosystem II can be transformed into free bicarbonate. Exogenous bicarbonate is a polar substance that cannot easily enter the oxygen-evolving center of the photosystem II under short saturating light flashes. Therefore, assuming that bicarbonate photolysis does occur in the photosystem II, this photosynthetic O_2 evolution should be elicited by bound bicarbonate rather than by exogenous bicarbonate (Fig. 3a). The evolved O_2 elicited by bound bicarbonate is of course ordinary oxygen, and the amount of oxygen evolution is also one order of magnitude less than that elicited by exogenous bicarbonate.

The experiment also showed that the labeled O_2 signal increased 8-fold compared to the unlabeled signal when 100 μl heavy oxygen or 3 μl ordinary water was respectively added in total darkness to dark-adapted normal chloroplasts under short saturating light flashes. This does not bring sole explanation that the oxygen flash yielded comes only from the water. Water does not easily enter the cell membrane, and the most common way for water to pass through chloroplast membrane is simple and slow diffusion. However, there are many CAs in the chloroplast, even in the thylakoids. The soluble CA in the stroma works in coordination with the intrinsic and extrinsic CAs in the thylakoids to form an inorganic carbon pool in the thylakoids, where the rapid group exchange of oxygen in H_2O and inorganic carbon occur. The rate of diffusion on the chloroplast membrane of 100 μl heavy oxygen water was one order of magnitude greater than that of 3 μl ordinary water, and the amount of bicarbonate transformed by heavy oxygen water and CO_2 was also one order of magnitude greater than that transformed by ordinary water and CO_2 in the thylakoids. Thus, the labeled O_2 signal elicited by bicarbonate transformed from 0.1 ml H_2^{18}O is one order of magnitude greater than the unlabeled O_2 signal elicited by bicarbonate transformed from 3 μl of ordinary water in the photosystem II after bicarbonate photolysis under the action of short saturating light flashes.

4.4 Revisiting the heavy oxygen (O^{18}) labeling experiment from Hillier et al. (2006)

The experiment was performed using spinach photosystem II membrane fragments illuminated by 16 saturating light flashes (5 Hz repetition rate) in the presence of 10 μM ethoxysolamide. Hillier et al. found that almost all the evolved oxygen was ordinary oxygen in the presence of labeled or unlabeled bicarbonate (Hillier et al. 2006). However, the above result cannot draw the conclusion on that bicarbonate is not a physiologically significant substrate. The isotopic exchange of oxygen among bicarbonate, CO_2 and the solvent water, and the conversion of bicarbonate to CO_2 and H_2O were significantly inhibited

(a)



(b)





Fig. 3 Bicarbonate photolysis was performed to explain two classic heavy oxygen labeling experiments under flashes. (a) In the experiment from Radmer and Ollinger (Radmer and Ollinge 1980), the oxygen evolved by HCO_3^- -depleted chloroplasts under short saturating light flashes comes from bound bicarbonate, and it is independent of exogenous HCO_3^- used for activation. (b) In the experiment from Hillier et al. (Hillier et al. 2006), most bicarbonates in photosystem II that were produced via the conversion of atmospheric CO_2 and water as solvent are unlabeled. Moreover, the majority of the bicarbonates used for photolysis are not labeled

when ethoxzolamide is used to inhibit the activity of intrinsic CAs in photosystem II. However, atmospheric CO_2 is not removed from the reaction system. Before the 16 flashes, labeled or unlabeled bicarbonate was injected into the reaction system. Obviously, less CO_2 in the reaction system comes from the added bicarbonate as ethoxzolamide inhibits the activity of the intrinsic CAs in

photosystem II. In addition, water-soluble bicarbonate cannot easily enter, however, liposoluble CO_2 can easily enter the photosystem II membrane, where extremely rapid CO_2 hydration can take place to form bicarbonate (Lu and Stemler 2007), with most bicarbonate in photosystem II being produced via the conversion of water as solvent and atmospheric CO_2 . Therefore, if bicarbonate photolysis in

Fig. 4 The combined effort of bicarbonate and water in photosynthetic oxygen evolution. The premise of water photolysis is bicarbonate photolysis. Bicarbonate photolysis and water photolysis work together in a 1:1 (mol/mol) stoichiometric ratio, releasing oxygen and carbon dioxide in a 1:1 (mol/mol) stoichiometry



photosystem II produces oxygen, the oxygen evolved is mostly ordinary oxygen (Fig. 3b).

5 Bicarbonate photolysis and water photolysis as well as their roles in photosynthesis

If photosynthetic organisms only perform photosynthetic oxygen evolution through bicarbonate photolysis, the following reaction occurs: $H_2O + CO_2 \rightarrow H^+ + HCO_3^- \rightarrow 1/2 O_2 + 2e^- + 2 H^+ + CO_2$ (Wu 2021a). It can be seen from the reaction equation that 2 moles of bicarbonate provide 1 mol of oxygen and 2 moles of CO_2 . This would mean assimilation is twice that of dissimilation to maintain the metabolism balance in nature, which seems unreasonable. If photosynthetic oxygen evolution produced by photosynthetic organisms occurs only through water photolysis, the following reaction occurs in photosynthetic organisms: $H_2O \rightarrow 1/2 O_2 + 2e^- + 2 H^+$. It can be seen from the reaction equation that 2 moles of water can provide only 1 mol of oxygen and there is no stoichiometric relationship between the water photolysis and the carbon dioxide assimilation (Fig. 1). Therefore, both bicarbonate photolysis and water photolysis account for half of the photosynthetic oxygen evolution respectively, which can not

only explain bicarbonate effect and carbon dioxide concentrating mechanisms (CCMS) in photosystem II, but also elaborate the origin of oxygen evolved in overall chemical reaction of photosynthesis more clearly than the traditional explanation (Fig. 4). The synthetic formula of photosynthetic oxygen evolution can be written as follows: $2H_2O + CO_2 \rightarrow H_2O + H^+ + HCO_3^- \rightarrow O_2 + 4e^- + 4 H^+ + CO_2$. It can be seen from the synthetic formula that photosynthetic oxygen evolution occurs in the presence of inorganic carbon. In other words, bicarbonate photolysis governs photosynthetic oxygen evolution (Fig. 4).

Bicarbonate photolysis is the premise of water photolysis, which is accompanied by bicarbonate photolysis during the photosynthetic oxygen evolution; this may explain why neither water photolysis nor bicarbonate photolysis is performed by anaerobic photosynthetic bacteria such as sulfur bacteria, in the presence of both inorganic carbon and water. These bacteria live in an acidic environment where inorganic carbon exists in the form of carbon dioxide rather than bicarbonate, and thus they do not undergo bicarbonate photolysis; accordingly, no water photolysis under light (Cohen et al. 1975). These bacteria only have acidic substances such as hydrogen sulfide photolysis, which leads to their photosynthesis without oxygen evolution.

It can be seen from the reaction equation of bicarbonate photolysis that CO₂ obviously acts as a concurrent post of “catalyst” to accelerate the photosynthetic oxygen evolution in photosynthetic cells (bicarbonate effect), which was called “activated CO₂” by Warburg and Krippahl (Warburg and Krippahl 1958). The bicarbonate-mediated CO₂ formation closely correlating with O₂ evolution were found on both the donor and acceptor side of PSII, which confirm the role of CO₂ as a concurrent “catalyst” and authenticity of the reaction equation of bicarbonate photolysis (Shevela et al. 2020).

The existence of Dole effect is in serious conflict with the view that the oxygen in atmosphere only comes from water photolysis by photosynthetic organism. However, we calculated the ¹⁸O enrichment in atmospheric O₂ according to that bicarbonate photolysis and water photolysis account for half the photosynthetic oxygen evolution respectively, and found that ¹⁸O enrichment in atmospheric O₂ with respect to water and bicarbonate (0.2043) is infinite approach to actual ¹⁸O content in the atmosphere (0.2041) (using (Metzner 1975) data to calculate). It demonstrated that the oxygen in the atmosphere sourced undoubtedly from water and bicarbonate evenly, and the Dole effect does not exist when ¹⁸O enrichment in atmospheric O₂ with respect to water and bicarbonate was considered.

Meanwhile, bicarbonate photolysis can not only release oxygen but also provide CO₂ for ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco), elevate and concentrate the CO₂ concentration in Rubisco, accelerate the absorption and reduction of inorganic carbon, and suppress the oxidation of organic matter. Carbon dioxide concentrating mechanisms (CCMs) in photosystem II, which are responsible for CO₂ production via bicarbonate photolysis, tend to be, major, primitive, indispensable and crucial for plant survival and growth. CCMs in the plasma membrane function as subordinate (Wu 2021b).

Last but not least, from the entire process of photosynthetic oxygen evolution, it can be seen that the supply of inorganic carbon determines the water demand of plants, which coordinates the balance of water metabolism and carbon metabolism, and ensures that plants to adapt to the environment. For example, when C₃ plants were subjected to drought stress, their stomata closed. Accordingly, their carbon dioxide supply has stopped, which terminates water photolysis and bicarbonate photolysis, and eliminates the demand for water simultaneously. When the stomata opening increases with a rise in carbon dioxide supply, water photolysis and bicarbonate photolysis also increase. In addition, when plants are in water deficit, they can boost the utilization of root-derived bicarbonate, which can greatly compensate for the shortage of water supply from the environment (Rao and Wu 2017).

6 Synergistic evolution of Earth spheres and carbon neutrality in nature

The Earth’s ecosystem can manage its own activities and maintain its own substance balance. Carbon and water balance play key roles in the Earth’s ecosystem. If oxygen evolved only from water photolysis in photosynthetic organisms on Earth in the early stage, it would be difficult for photosynthetic organisms to couple with the hydrosphere and the atmosphere, and it would be even more difficult to couple with the lithosphere (Fig. 1). However, the photosynthetic oxygen evolved from bicarbonate photolysis and water photolysis can stoichiometrically couple with the hydrosphere and atmosphere of the early Earth and then couple with the lithosphere, thereby realizing carbon and water balance (Wu 2022) (Fig. 5).

From the synergistic effects (changes) of CO₂ and H₂O in the atmosphere, bicarbonate, CO₂ partial pressure and pH in the hydrosphere, and carbonate rocks in the lithosphere on Earth in the early stage, photosynthetic oxygen evolution is actually stoichiometrically coupled with the atmosphere, hydrosphere and lithosphere (Hart 1978; Kasting 1993; Des Marais 2000; Dismukes et al. 2001; Halevy and Bachan 2017; Wu 2022) (Fig. 5).

Furthermore, the photosynthetic oxygen evolved from bicarbonate photolysis and water photolysis represents the coupling as 1:1 (mol/mol) stoichiometric relationship between the reduction of CO₂ and oxidation of organic carbon, as well as between the substance cycle and energy cycle in the biosphere, which contributes to “carbon neutrality” in the whole Earth system (Fig. 6). Introducing a more accurate process of photosynthetic oxygen evolution into textbooks can assist future generations in learning from nature in managing carbon balance and water balance.

7 Conclusion

Many types of CA have been found in chloroplasts. Meanwhile, photosystem II functions as CA to catalyze the reaction of CO₂ hydration under physiological conditions, which leads to almost complete isotope exchange between oxygen in water and oxygen in inorganic carbon in the Hill reaction system. This recent knowledge and research progress enable us to draw new conclusions based on bicarbonate effect, Dole effect, the characteristics of photosystem II, and several classic heavy oxygen labeling experiments. Photosynthetic oxygen evolution should source from both bicarbonate and water as 1:1 (mol/mol) stoichiometric relationship, which initiates the journey of water metabolism and carbon metabolism in nature,




Fig. 5 The coupling of photosynthetic oxygen evolution and Earth spheres. Carbonate rocks (lithosphere) are dissolved under the action of water (hydrosphere) and carbon dioxide (atmosphere) to form Ca²⁺ and bicarbonate (biosphere). Ca²⁺ combines with bicarbonate to form the precipitation of calcium carbonate (CaCO₃) into the lithosphere. Plants split bicarbonate and water to release oxygen and carbon dioxide (biosphere). Plants then assimilate carbon dioxide to form carbohydrates (CH₂O) (biosphere). Finally, organisms utilize oxygen to decompose carbohydrates into carbon dioxide to the atmosphere and water to the hydrosphere




Fig. 6 Carbon neutrality and water neutrality in nature driven by photosynthetic oxygen evolution. The photosynthetic oxygen evolution, which converts light energy into chemical energy, is performed by the synergistic action of bicarbonate photolysis (①) and water photolysis (②) with a stoichiometric relationship of 1:1 (mol/mol). It produces oxygen and CO₂ in a stoichiometric ratio of 1:1, which determines that inorganic carbon assimilation (③) is equal to organic carbon dissimilation (④). Finally, from the starting point (photosynthetic oxygen evolution) to the endpoint (carbon dissimilation) of the carbon neutrality strategy, the net flux of both CO₂ and water in nature is 0

solidifies the relationship of assimilation equal to dissimilation, and provides a demonstration of how to coordinate water and carbon balances to maintain water neutrality and carbon neutrality (Fig. 6). In the future, more direct evidence about that the O₂ evolution in photosynthesis evenly comes from water and bicarbonate will need to provide.

Acknowledgements We thank the foundations of the National Natural Science Foundation of China [No. U1612441-2], and Support Plan Projects of Science and Technology Department of Guizhou Province [No. (2021)YB453].

Declarations

Conflict of interest The author declares no conflict of interest.

References

- Allen FL (1955) Photosynthetic evolution of oxygen by flashes of light. *Arch Biochem Biophys* 58(1):124–143
- Bender M, Sowers T, Labeyrie L (1994) The dole effect and its variations during the last 130,000 years as measured in the Vostok ice core. *Global Biogeochem Cycles* 8(3):363–376
- Bowden SJ, Hallahan BJ, Ruffle SV, Evans MCW, Nugent JHA (1991) Preparation and characterization of Photosystem II core particles with and without bound bicarbonate. *BBA-Bioenergetics* 1060(1):89–96
- Cohen Y, Jørgensen BB, Padan ET, Shilo M (1975) Sulphide-dependent anoxygenic photosynthesis in the cyanobacterium *Oscillatoria limnetica*. *Nature* 257(5526):489–492
- Dai X, Yu Y, Zhang R, Yu X, He P, Xu C (2001) Relationship among photosystem II carbonic anhydrase, extrinsic polypeptides and manganese cluster. *Chin Sci Bull* 46(5):406–408
- Delosme R, Joliot P (2002) Period four oscillations in chlorophyll a fluorescence. *Photosynth Res* 73:165–168
- Des Marais DJ (2000) When did photosynthesis emerge on Earth? *Science* 289(5485):1703–1705
- Dismukes GC, Klimov VV, Baranov SV, Kozlov YN, DasGupta J, Tyryshkin A (2001) The origin of atmospheric oxygen on Earth: the innovation of oxygenic photosynthesis. *P Natl Acad Sci USA* 98(5):2170–2175
- Ferreira KN (2004) Architecture of the photosynthetic oxygen-evolving center. *Science* 303(5665):1831–1838
- Govindjee, Beatty JT, Gest H, Allen JF (2006) Discoveries in photosynthesis. Springer, Dordrecht, pp 63–105
- Halevy I, Bachan A (2017) The geologic history of seawater pH. *Science* 355(6329):1069–1071
- Hart MH (1978) The evolution of the atmosphere of the Earth. *Icarus* 33(1):23–39
- Hill R (1937) Oxygen evolved by isolated chloroplasts. *Nature* 139(3525):881–882
- Hillier W, McConnell I, Badger MR, Boussac A, Klimov VV, Dismukes GC, Wydrzynski T (2006) Quantitative assessment of intrinsic carbonic anhydrase activity and the capacity for bicarbonate oxidation in photosystem II. *Biochemistry* 45(7):2094–2102
- Hoffmann G, Cuntz M, Weber C, Ciais P, Friedlingstein P, Heimann M, Jouzel J, Kaduk J, Maier-Reimer E, Seibt U, Six K (2004) A model of the Earth's dole effect. *Global Biogeochem Cycles* 8(1):18 (GB1008)
- Hogetsu D, Miyachi S (1979) Role of carbonic anhydrase in photosynthetic CO₂ fixation in *Chlorella*. *Plant Cell Physiol* 20(4):747–756
- Ignatova LK, Rudenko NN, Mudrik VA, Fedorchuk TP, Ivanov BN (2011) Carbonic anhydrase activity in *Arabidopsis thaliana* thylakoid membrane and fragments enriched with PSI or PSII. *Photosynth Res* 110:89–98
- Joliot P (2003) Period-four oscillations of the flash-induced oxygen formation in photosynthesis. *Photosynth Res* 76(1):65–72
- Kasting JF (1993) Earth's early atmosphere. *Science* 259(5097):920–926
- Khrustin MS, Ignatova LK, Rudenko NN, Ivanov BN, Klimov VV (2004) Photosystem II associated carbonic anhydrase activity in higher plants is situated in core complex. *FEBS Lett* 577:305–308
- Kok B, Forbusch B, McGloin M (1970) Cooperation of charges in photosynthetic O₂ evolution-I. A linear four step mechanism. *Photochem Photobiol* 11:457–475
- Koroidov S, Shevela D, Shutova T, Samuelsson G, Messinger J (2014) Mobile hydrogen carbonate acts as proton acceptor in photosynthetic water oxidation. *P Natl Acad Sci USA* 111(17):6299–6304
- Lu YK, Stemler AJ (2007) Differing responses of the two forms of photosystem II carbonic anhydrase to chloride, cations, and pH. *BBA-Bioenergetics* 1767:633–638
- Mader M, Schmidt C, van Geldern R, Barth JA (2017) Dissolved oxygen in water and its stable isotope effects: a review. *Chem Geol* 473:10–21
- Metzner H (1975) Water decomposition in photosynthesis? A critical reconsideration. *J Theor Bio* 51(1):201–231
- Moskvin OV, Shutova TV, Khrustin MS, Ignatova LK, Villarejo A, Samuelsson G, Ivanov BN (2004) Carbonic anhydrase activities in pea thylakoids. *Photosynth Res* 79(1):93–100
- Moubarak-Milad M, Stemler A (1994) Oxidation-reduction potential dependence of photosystem II carbonic anhydrase in maize thylakoids. *Biochemistry* 33(14):4432–4438
- Okrasav K, Kazlauskas RJ (2006) Manganese-substituted carbonic anhydrase as a new peroxidase. *Chem-Eur J* 12(6):1587–1596
- Pronina NA, Allakhverdiev SI, Kupriyanova EV, Klyachko-Gurvich GL, Klimov VV (2002) Carbonic anhydrase in subchloroplast particles of pea plants. *Russ J Plant Phys* 49(3):303–310
- Radmer R, Ollinger O (1980) Isotopic composition of photosynthetic O₂ flash yields in the presence of H₂¹⁸O and HC¹⁸O⁻. *FEBS Lett* 110(1):57–61
- Rao S, Wu Y (2017) Root-derived bicarbonate assimilation in response to variable water deficit in *Camptotheca acuminata* seedlings. *Photosynth Res* 134(1):59–70
- Rodionova MV, Zhamukhamedov SK, Karacan MS, Venedik KB, Allakhverdiev SI (2017) Evaluation of new cu(II) complexes as a novel class of inhibitors against plant carbonic anhydrase, glutathione reductase, and photosynthetic activity in photosystem II. *Photosynth Res* 133(1):1–15
- Ruben S, Randall M, Kamen M, Hyde JL (1941) Heavy oxygen (O¹⁸) as a tracer in the study of photosynthesis. *J Am Chem Soc* 63(3):877–879
- Rudenko NN, Ignatova LK, Kamornitskaya VB, Ivanov BN (2006) Pea leaf thylakoids contain several carbonic anhydrases. *Dokl Biochem Biophys* 408(1):155–157
- Rudenko NN, Ignatova LK, Ivanov BN (2007) Multiple sources of carbonic anhydrase activity in pea thylakoids: soluble and membrane-bound forms. *Photosynth Res* 91(1):81–89
- Rudenko NN, Ignatova LK, Fedorchuk TP, Ivanov BN (2015) Carbonic anhydrases in photosynthetic cells of higher plants. *Biochem Moscow* 80(6):674–687

- Shevela D, Eaton-Rye JJ, Shen JR, Govindjee (2012) Photosystem II and the unique role of bicarbonate: a historical perspective. *BBA-Bioenergetics* 1817(8):1134–1151
- Shevela D, Do HN, Fantuzzi A, Rutherford AW, Messinger J (2020) Bicarbonate-mediated CO₂ formation on both sides of photosystem II. *Biochemistry* 59(26):2442–2449
- Shitov AV, Zharmukhamedov SK, Shutova TV, Allakhverdov SI, Samuelsson G, Klimov VV (2011) A carbonic anhydrase inhibitor induces bicarbonate-reversible suppression of electron transfer in pea photosystem 2 membrane fragments. *J Photoch Photobio B* 104(1–2):366–371
- Shitov AV, Terentyev VV, Zharmukhamedov SK, Rodionova MV, Karacan M, Karacan N, Klimov VV, Allakhverdov SI (2018) Is carbonic anhydrase activity of photosystem II required for its maximum electron transport rate? *BBA Ioenergetics* 1859(4):292–299
- Stemler A (1980) Inhibition of photosystem II by formate: possible evidence for a direct role of bicarbonate in photosynthetic oxygen evolution. *BBA-Bioenergetics* 593:103–112
- Stemler AJ (2002) The bicarbonate effect, oxygen evolution and the shadow of Otto Warburg. *Photosynth Res* 73:177–183
- Stemler A, Radmer R (1975) Source of photosynthetic oxygen in bicarbonate-stimulated Hill reaction. *Science* 190(4213):457–458
- Tikhonov K, Shevela D, Klimov VV, Messinger J (2018) Quantification of bound bicarbonate in photosystem II. *Photosynthetica* 56(1):210–216
- Umena Y, Kawakami K, Shen JR, Kamiya N (2011) Crystal structure of oxygen-evolving photosystem α - at a resolution of 1.9 Å. *Nature* 473(7345):55–60
- Van Rensen JJS, Vermaas WFJ (1981) Action of bicarbonate and photosystem II inhibiting herbicides on electron transport in pea grana and in thylakoids of a blue-green alga. *Physiol Plant* 51:106–110
- Van Rensen JJS, Xu C, Govindjee (1999) Role of bicarbonate in photosystem II, the water-plastoquinone oxido-reductase of plant photosynthesis. *Physiol Plant* 105(3):585–592
- Warburg O, Krippahl G (1958) Hill-reaktionen. *Z Naturforsch B* 13(8):509–514
- Warburg O, Krippahl G, Jetschmann C (1965) Widerlegung der photolyse des wassers und beweis der photolyse der kohlensäure nach versuchen mit lebender *Chlorella* und den Hill-reagentien nitrat und K₃Fe(CN)₆. *Z Naturforsch B* 20(10):993–996
- Wu Y (2021a) Is bicarbonate directly used as substrate to participate in photosynthetic oxygen evolution. *Acta Geochim* 40(4):650–658
- Wu Y (2021b) Bicarbonate use and carbon dioxide concentrating mechanisms in photosynthetic organisms. *Acta Geochim* 40(5):846–853
- Wu Y (2022) The increase in the karstification–photosynthesis coupled carbon sink and its implication for carbon neutrality. *Agronomy* 12:2147

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.