

# Effects of carbon anhydrase on utilization of bicarbonate in microalgae: a case study in Lake Hongfeng

Haitao Li<sup>1,2</sup> · Yanyou Wu<sup>1</sup> · Lihua Zhao<sup>1</sup>

Received: 25 November 2017/Revised: 23 April 2018/Accepted: 18 May 2018/Published online: 1 June 2018  
© Science Press, Institute of Geochemistry, CAS and Springer-Verlag GmbH Germany, part of Springer Nature 2018

**Abstract** A bidirectional labeling method was established to distinguish the proportions of  $\text{HCO}_3^-$  and  $\text{CO}_2$  utilization pathways of microalgae in Lake Hongfeng. The method was based on microalgae cultured in a medium by adding equal concentrations of  $\text{NaH}^{13}\text{CO}_3$  with different  $\delta^{13}\text{C}$  values simultaneously. The inorganic carbon sources were quantified according to the stable carbon isotope composition in the treated microalgae. The effects of extracellular carbonic anhydrase (CAex) on the  $\text{HCO}_3^-$  and  $\text{CO}_2$  utilization pathways were distinguished using acetazolamide, a potent membrane-impermeable carbonic anhydrase inhibitor. The results show utilization of the added  $\text{HCO}_3^-$  was only 8% of the total carbon sources in karst lake. The proportion of the  $\text{HCO}_3^-$  utilization pathway was 52% of total inorganic carbon assimilation. Therefore, in the natural water of the karst area, the microalgae used less bicarbonate that preexisted in the aqueous medium than  $\text{CO}_2$  derived from the atmosphere. CAex increased the utilization of inorganic carbon from the atmosphere. The microalgae with CAex had greater carbon sequestration capacity in this karst area.

**Keywords** Microalgae · Carbonic anhydrase · Stable carbon isotope · Inorganic carbon utilization

## 1 Introduction

High pH and high concentration of bicarbonate are two typical characteristics of karst lakes. The main component of karst is  $\text{MgCa}(\text{CO}_3)_2$ , a highly soluble rock. The proportion of  $\text{CO}_2$  in total dissolved inorganic carbon (DIC) is less than 1% in high pH conditions (Riebesell et al. 1993). Thus, the  $\text{CO}_2$  in aquatic media that can be directly utilized by photosynthesis in microalgae is limited (Talling 1976). Several microalgae have adapted by forming carbon-concentrating mechanisms (CCMs) to increase  $\text{CO}_2$  concentrations to meet their photosynthetic demands (Colman et al. 2002; Giordano et al. 2005). Another strategy is to utilize bicarbonate (Colman et al. 2002). Carbonic anhydrase (CA) may play the key role in these carbon assimilation systems.

CA (EC 4.2.1.1), a zinc-containing metalloenzyme, catalyzes the reversible interconversion between  $\text{HCO}_3^-$  and  $\text{CO}_2$ . CA is one of the most important enzymes in physiological processes and significantly accelerates the photosynthetic assimilation of inorganic carbon (Ci) (Badger and Price 1994; Sültemeyer 1998). CA is widely distributed and multiple types exist in microalgae. One of the most important CAs is extracellular CA (CAex), which may be involved in CCMs and in  $\text{HCO}_3^-$  utilization (Williams and Turpin 1987; Badger and Price 1994; Elzenga et al. 2000; Mondal et al. 2016).

Stable carbon isotope ( $\delta^{13}\text{C}$ ) analysis is an important tool to identify various Ci sources (Fry and Sherr 1984; Bade et al. 2006; Chen et al. 2009). Different Ci sources and assimilation mechanisms cause variations in  $\delta^{13}\text{C}$  fractionation. The  $\text{HCO}_3^-$  in the uncatalyzed pathway produces approximately 10‰ of  $\delta^{13}\text{C}$  fractionation (Mook et al. 1974), while  $\text{HCO}_3^-$  assimilation catalyzed by CAex produces only 1.1‰ of  $\delta^{13}\text{C}$  fractionation (Marlier and

✉ Yanyou Wu  
wuyanyou@vip.skleg.cn

<sup>1</sup> State Key Laboratory of Environmental Geochemistry/  
Institute of Geochemistry, Chinese Academy of Sciences,  
Guiyang 550081, China

<sup>2</sup> School of Karst Science, Guizhou Normal University/State  
Engineering Technology Institute for Karst Desertification  
Control, Guiyang 550001, China

O'Leary 1984). An approximately 9‰ discrimination of carbon isotope has been found between the  $\text{HCO}_3^-$  catalyzed by CAex and that uncatalyzed (Wu et al. 2012).

Acetazolamide (AZ) is a potent membrane-impermeable CA inhibitor that selectively inhibits CAex activity (Moroney et al. 1985). The addition of AZ enables determination of the effect of CAex on Ci utilization.

Several studies have investigated mechanisms of Ci utilization in microalgal species (Axelsson et al. 1995; Mozami-Goudarzi and Colman 2011; Moulin et al. 2011; Smith-Harding et al. 2017). However, the conventional technique cannot quantify the proportions of DIC sources and their microalgal pathways in karst lakes (Xie and Wu 2017). This is the aim of this study.

To this end, microalgae from Lake Hongfeng were cultured in different concentrations of  $\text{NaHCO}_3$  and AZ. The proportion of Ci sources and pathways were determined by comparing their  $\delta^{13}\text{C}$  compositions under separate experiments adding two labeled  $\delta^{13}\text{C}$  bicarbonates. We then estimated the contribution of microalgal CAex to Ci sources and utilization pathways in the karst lake.

## 2 Materials and methods

### 2.1 Research site

Lake Hongfeng (106°19' to 106°28'E, 26°26' to 26°35'N) is in central Guizhou Province in the core of the southwest karst area of China. The concentration of  $\text{HCO}_3^-$  in Lake Hongfeng is 1.0–2.5 mmol/L, and the pH is  $8.1 \pm 0.4$  (Wu et al. 2008).

### 2.2 Microalgae incubation

The microalgal samples were obtained from Lake Hongfeng. All samples were incubated at  $25.0 \pm 1.0$  °C under a  $150 \mu\text{mol m}^{-2} \text{s}^{-1}$  light intensity with a 12/12 h day/night cycle. The pH was adjusted to 8.10 by adding NaOH. AZ was obtained from Sigma-Aldrich Co. (St. Louis, USA).

The microalgae were grown in the lake water in Erlenmeyer flasks after filtration through a 45- $\mu\text{m}$  glass microfiber filter. Treatments are listed in Table 1. The cultures were treated for 5 days.

### 2.3 Measurement of the microalgal growth

Microalgal protein was analyzed using the method of Coomassie Brilliant Blue (Sedmak and Grossberg 1977). A volume of 5–10 ml of microalgae was centrifuged, and then resuspended. The optical density was tested using a spectrophotometer (Labtech UV-2000, Boston, USA) at

**Table 1** Experimental treatments of the microalgae

Treatment	$\text{NaHCO}_3$ (mmol/L)	AZ (mmol/L)	$\delta^{13}\text{C}_{\text{bicarbonate}}$
1	1.0	0, 1.0, 10.0	①, ②
2	2.5	0, 1.0, 10.0	①, ②
3	5.0	0, 1.0, 10.0	①, ②
4	20.0	0, 1.0, 10.0	①, ②

①: cultured in  $\text{NaHCO}_3$  with the  $\delta^{13}\text{C}$  value of  $-17.4\text{‰}$ (‰, PDB)

②: cultured in  $\text{NaHCO}_3$  with the  $\delta^{13}\text{C}$  value of  $-28.4\text{‰}$ (‰, PDB)

595 nm (OD595). The protein content is expressed as  $\mu\text{g/L}$  based on the aqueous medium.

### 2.4 Measurement of $\delta^{13}\text{C}$ in microalgae

The microalgae materials were freeze-dried and then converted to  $\text{CO}_2$  at 850 °C in a quartz tube with copper oxide to provide oxygen for combustion. The extracted  $\text{CO}_2$  from the samples was purified as follows.

Water and oxygen were removed from the gas stream using two traps. The first was an alcohol–liquid nitrogen mixture to separate the water vapor, and the second was liquid nitrogen to condense the  $\text{CO}_2$ . After this double distillation, the isolated  $\text{CO}_2$  was collected into a sample tube. The  $\text{CO}_2$  sample was analyzed with an isotope ratio mass spectrometer (Finnigan MAT 252, Bremen, Germany). All isotopic compositions ( $\delta^{13}\text{C}$ ) are expressed as per mille (‰) and compared with the Pee Dee Belemnite (PDB) standard [see Eq. (1)]. The analytical precision was  $\pm 0.1\text{‰}$ .

$$\delta^{13}\text{C} (\text{‰}) = [(R_{\text{sample}}/R_{\text{standard}}) - 1] \times 1000 \quad (1)$$

where  $R_{\text{sample}}$  and  $R_{\text{standard}}$  are the ratios of heavy to light isotope ( $^{13}\text{C}/^{12}\text{C}$ ) of the sample and the standard, respectively.

### 2.5 Distinguishing the different carbon sources and metabolic pathways

This study chose an open system to simulate natural conditions. In this type of system, Ci in the liquid medium and the atmosphere is in dynamic balance. We created a bidirectional labeling method that simultaneously cultured microalgae in  $\text{NaHCO}_3$  of different  $\delta^{13}\text{C}$  to address this problem.

There is an assumption that the proportion of added Ci utilization is the same under the same concentration of  $\text{HCO}_3^-$  at the same time regardless of which labeled  $\text{HCO}_3^-$  was added. This is the theoretical basis of the bidirectional labeling method.

In general, algae can utilize DIC from the atmosphere and added  $\text{HCO}_3^-$ . Therefore, the  $\delta^{13}\text{C}$  of the algae was fit for the bivariate isotope-mixture model that can be expressed as:

$$\delta_{\text{Ai}} = (1 - f_{\text{bi}})\delta_{\text{a}} + f_{\text{bi}}(\delta_{\text{a}} + 9\text{‰}) \quad (2)$$

where  $\delta_{\text{Ti}}$  is the  $\delta^{13}\text{C}$  value of the algae cultured in the same concentration of  $\text{HCO}_3^-$  with different  $\delta^{13}\text{C}$  values;  $f_{\text{Bi}}$  is the proportion of the utilization of DIC from the added  $\text{HCO}_3^-$  in the total carbon sources used by the microalgae; and  $\delta_{\text{Ai}}$  and  $\delta_{\text{Bi}}$  are the  $\delta^{13}\text{C}$  values of the algae after using DIC from atmospheric  $\text{CO}_2$  or from the added  $\text{HCO}_3^-$ , respectively, as their sole carbon sources.

In this experiment, the microalgae could utilize both  $\text{CO}_2$  and  $\text{HCO}_3^-$  as carbon sources. Approximately 9‰ carbon isotope discrimination has been observed between  $\text{CO}_2$  and  $\text{HCO}_3^-$  utilization pathways (Wu et al. 2012). Therefore,  $\delta_{\text{Ai}}$  and  $\delta_{\text{Bi}}$  can be expressed as follows:

$$\delta_{\text{Ai}} = (1 - f_{\text{bi}})\delta_{\text{a}} + f_{\text{bi}}(\delta_{\text{a}} + 9\text{‰}) \quad (3)$$

$$\delta_{\text{Bi}} = (1 - f_{\text{bi}})\delta_{\text{ai}} + f_{\text{bi}}(\delta_{\text{ai}} + 9\text{‰}) \quad (4)$$

where  $f_{\text{bi}}$  is the proportion of the  $\text{HCO}_3^-$  pathway;  $\delta_{\text{a}}$  is the  $\delta^{13}\text{C}$  value of the algae after using DIC in the sole form of the  $\text{CO}_2$  utilization pathway from the atmospheric source; and  $\delta_{\text{ai}}$  is the  $\delta^{13}\text{C}$  value of the algae after using DIC in the sole form of the  $\text{CO}_2$  utilization pathway of the added  $\text{HCO}_3^-$  source.

Based on Eqs. (3) and (4), Eq. (2) can be expressed as:

$$\delta_{\text{Ti}} = (1 - f_{\text{Bi}})[(1 - f_{\text{bi}})\delta_{\text{a}} + f_{\text{bi}}(\delta_{\text{a}} + 9\text{‰})] + f_{\text{Bi}}[(1 - f_{\text{bi}})\delta_{\text{ai}} + f_{\text{bi}}(\delta_{\text{ai}} + 9\text{‰})] \quad (i = 1, 2) \quad (5)$$

Equation (5) can be simplified to:

$$\delta_{\text{Ti}} = \delta_{\text{a}} + f_{\text{Bi}}(\delta_{\text{ai}} - \delta_{\text{a}}) + 9\text{‰}f_{\text{bi}} \quad (i = 1, 2) \quad (6)$$

For the two labeled types of  $\text{NaHCO}_3$  in our experiments, Eq. (6) can be rewritten as:

$$\delta_{\text{T1}} = \delta_{\text{a}} + f_{\text{B1}}(\delta_{\text{a1}} - \delta_{\text{a}}) + 9\text{‰}f_{\text{b1}} \quad (7)$$

$$\delta_{\text{T2}} = \delta_{\text{a}} + f_{\text{B2}}(\delta_{\text{a2}} - \delta_{\text{a}}) + 9\text{‰}f_{\text{b2}} \quad (8)$$

There are some important facts that we should note in Eqs. (7) and (8). The first is that the proportion of added Ci utilization is the same at the same concentration of  $\text{HCO}_3^-$  regardless of which labeled  $\text{NaHCO}_3$  was added. Therefore,  $f_{\text{B1}} = f_{\text{B2}} = f_{\text{B}}$ . The second is that the proportion of the  $\text{HCO}_3^-$  pathway is the same at the same concentration of  $\text{HCO}_3^-$  regardless of the labeled  $\delta^{13}\text{C}$  of  $\text{NaHCO}_3$ . Therefore,  $f_{\text{b1}} = f_{\text{b2}} = f_{\text{b}}$ . Although  $\delta_{\text{a1}}$  and  $\delta_{\text{a2}}$  cannot be obtained, the difference between them is simply the difference between  $\delta^{13}\text{C}$  values of the first and second labeled  $\text{NaHCO}_3$  in the medium. Therefore, the  $(\delta_{\text{a1}} - \delta_{\text{a2}})$  can be

replaced with  $(\delta_{\text{C1}} - \delta_{\text{C2}})$ . Therefore,  $f_{\text{B}}$  can be expressed as follows:

$$f_{\text{B}} = \frac{\delta_{\text{T1}} - \delta_{\text{T2}}}{\delta_{\text{C1}} - \delta_{\text{C2}}} \quad (9)$$

where  $\delta_{\text{C1}}$  and  $\delta_{\text{C2}}$  are the  $\delta^{13}\text{C}$  values of the first and second labeled  $\text{NaHCO}_3$  in the medium, respectively. From Eq. (9), it can be concluded that the proportion of the utilization of DIC from the added  $\text{HCO}_3^-$  in the total carbon sources used by the microalgae ( $f_{\text{B}}$ ) was dependent only on the  $\delta^{13}\text{C}$  values of the algae harvested and the labeled  $\text{NaHCO}_3$  added, regardless of DIC form and origin.

The  $(\delta_{\text{ai}} - \delta_{\text{a}})$  in Eq. (6) can be replaced with  $(\delta_{\text{Ci}} - \delta_{\text{C0}})$ . A new equation can then be formulated as:

$$\delta_{\text{ai}} - \delta_{\text{a}} = \delta_{\text{Ci}} - \delta_{\text{C0}} = D_i \quad (i = 1, 2, 3, \dots, n) \quad (10)$$

Therefore, Eq. (6) can be rewritten as:

$$\delta_{\text{Ti}} = \delta_{\text{a}} + f_{\text{Bi}}D_i + 9\text{‰}f_{\text{bi}} \quad (i = 1, 2, 3, \dots, n) \quad (11)$$

The proportion of the  $\text{HCO}_3^-$  pathway ( $f_{\text{b}}$ ) can then be calculated as:

$$f_{\text{bi}} = 1000(\delta_{\text{Ti}} - \delta_{\text{a}} - f_{\text{Bi}}D_i)/9 \quad (i = 1, 2) \quad (12)$$

When  $f_{\text{bi}} = 0$ , Eq. (12) can be rewritten as:

$$\delta_{\text{a}} = \delta_{\text{Ti}} - f_{\text{Bi}}D_i \quad (i = 1, 2) \quad (13)$$

From Eqs. (9), (12), and (13), the proportion of the  $\text{HCO}_3^-$  pathway by the microalgae ( $f_{\text{b}}$ ) can be calculated. It was dependent only on the  $\delta^{13}\text{C}$  values of the algae harvested and the labeled  $\text{NaHCO}_3$  added.

To analyze the complete picture of Ci utilization, the bidirectional labeling method ( $\text{NaH}^{13}\text{CO}_3$  with different  $\delta^{13}\text{C}$  values added) can solve the difficulties of the time course of parameters (concentrations and isotopic data in incubation).

## 2.6 Statistical analysis

All experiments were conducted in triplicate. Data are expressed as mean  $\pm$  standard error.

## 3 Results

### 3.1 Microalgae biomass

The content of protein in the treated microalgae increased in parallel with the  $\text{NaHCO}_3$  added (Table 2). However, it decreased with increasing concentrations of AZ added. Under the same concentration of  $\text{NaHCO}_3$ , microalgae growth was severely restricted by AZ. Compared to the control, the average effect of AZ on the microalgal protein content was 69% at 1.0 mmol/L AZ and 35% at

**Table 2** The protein contents( $\mu\text{g/L}$ ) and percentages of microalgae under bicarbonate and AZ treatments

[NaHCO <sub>3</sub> ][AZ] <sup>a</sup> mmol/L	0	1.0	10.0
1.0	1003.12 $\pm$ 14.37(100%)	768.45 $\pm$ 37.07(77%)	325.32 $\pm$ 37.39(32%)
2.5	1079.54 $\pm$ 134.65(100%)	753.33 $\pm$ 73.04(70%)	395.05 $\pm$ 46.88(37%)
5.0	1286.88 $\pm$ 79.90(100%)	806.29 $\pm$ 25.75(63%)	442.04 $\pm$ 13.19(34%)
20.0	1342.24 $\pm$ 41.24(100%)	913.34 $\pm$ 37.93(68%)	518.18 $\pm$ 94.60(39%)

<sup>a</sup>NaHCO<sub>3</sub> concentration added in the culture medium

10.0 mmol/L AZ. Among all treatments, the maximal growth rate treatment (20.0 mmol/L NaHCO<sub>3</sub> without AZ) was 4.12-fold higher than the minimal treatment (1.0 mmol/L NaHCO<sub>3</sub> with 10.0 mmol/L AZ).

### 3.2 Stable carbon isotope composition of the microalgae

The  $\delta^{13}\text{C}$  of the DIC was  $-11.0\text{‰} \pm 0.4\text{‰}$  which is positive relative to the  $\delta^{13}\text{C}$  of the added NaHCO<sub>3</sub> ( $-17.4\text{‰}$  or  $-28.4\text{‰}$ ). In the end, the  $\delta^{13}\text{C}$  value of the microalgae decreased as the amount of NaHCO<sub>3</sub> increased; it also decreased as AZ increased (for the same concentration of NaHCO<sub>3</sub>) (Table 3). The  $\delta^{13}\text{C}$  of the microalgae cultured in the treatment was affected by the added NaHCO<sub>3</sub> with different  $\delta^{13}\text{C}$  values. In general, the more negative the NaH<sup>13</sup>CO<sub>3</sub> added, the more negative the  $\delta^{13}\text{C}$  of the microalgae harvested for the same concentration of NaHCO<sub>3</sub> added.

### 3.3 Variation in carbon sources during different concentrations of NaHCO<sub>3</sub> and acetazolamide

Based on Eq. (9), we calculated the proportion of the utilization of DIC from the added NaHCO<sub>3</sub> ( $f_B$ ). The  $f_B$  increased with increasing NaHCO<sub>3</sub> concentration whether AZ was present or not (Table 4). In the treatment at 20.0 mmol/L NaHCO<sub>3</sub>,  $f_B$  increased with increasing concentration of AZ added.

**Table 4** The proportion of the utilization of DIC from the added HCO<sub>3</sub><sup>-</sup> to the total carbon sources under bicarbonate and AZ treatments

[NaHCO <sub>3</sub> ][AZ] <sup>a</sup> mmol/L	0	1.0	10.0
1.0	0.06 $\pm$ 0.03	0.06 $\pm$ 0.04	0.03 $\pm$ 0.03
2.5	0.08 $\pm$ 0.02	0.08 $\pm$ 0.05	0.09 $\pm$ 0.03
5.0	0.09 $\pm$ 0.04	0.17 $\pm$ 0.05	0.12 $\pm$ 0.05
20.0	0.20 $\pm$ 0.06	0.35 $\pm$ 0.09	0.40 $\pm$ 0.09

<sup>a</sup>NaHCO<sub>3</sub> or AZ concentration added in the culture medium separatelyThe  $\delta^{13}\text{C}$  values of the two kinds of NaHCO<sub>3</sub> added are  $-17.4\text{‰}$  or  $-28.4\text{‰}$  separately( $\text{‰}$ , PDB)

### 3.4 Variation in carbon pathways under different concentrations of NaHCO<sub>3</sub> and acetazolamide

Based on Eq. (12), the proportion of the HCO<sub>3</sub><sup>-</sup> pathway ( $f_b$ ) in microalgae was calculated. It decreased with increasing NaHCO<sub>3</sub> concentration in the treatment without AZ (Table 5). The  $f_b$  also decreased with increasing AZ concentration at the same concentration of added NaHCO<sub>3</sub>. In the treatment with 10.0 mmol/L AZ,  $f_b$  values were all very small ( $f_b \leq 0.12$ ).

**Table 3**  $\delta^{13}\text{C}$  of the microalgae cultured under bicarbonate and AZ treatments

Treatment [NaHCO <sub>3</sub> ] <sup>a</sup> (mmol/L)	0 mmol/L AZ		1.0 mmol/L AZ		10.0 mmol/L AZ	
	$\delta_{T1}$	$\delta_{T2}$	$\delta_{T1}$	$\delta_{T2}$	$\delta_{T1}$	$\delta_{T2}$
1.0	$-30.4 \pm 0.3$	$-31.0 \pm 0.1$	$-31.7 \pm 0.2$	$-32.4 \pm 0.3$	$-34.8 \pm 0.1$	$-35.2 \pm 0.3$
2.5	$-30.7 \pm 0.1$	$-31.6 \pm 0.1$	$-32.2 \pm 0.3$	$-33.1 \pm 0.4$	$-35.1 \pm 0.2$	$-36.0 \pm 0.2$
5.0	$-32.7 \pm 0.2$	$-33.7 \pm 0.3$	$-33.2 \pm 0.2$	$-35.1 \pm 0.2$	$-34.6 \pm 0.4$	$-36.0 \pm 0.3$
20.0	$-34.5 \pm 0.3$	$-36.7 \pm 0.3$	$-35.9 \pm 0.4$	$-39.8 \pm 0.3$	$-37.4 \pm 0.2$	$-41.7 \pm 0.3$

<sup>a</sup>NaHCO<sub>3</sub> concentration added in the culture medium $\delta_{T1}$ : cultured in NaHCO<sub>3</sub> with the  $\delta^{13}\text{C}$  value of  $-17.4\text{‰}$ ( $\text{‰}$ , PDB) $\delta_{T2}$ : cultured in NaHCO<sub>3</sub> with the  $\delta^{13}\text{C}$  value of  $-28.4\text{‰}$ ( $\text{‰}$ , PDB)

**Table 5** The proportion of the  $\text{HCO}_3^-$  pathways under bicarbonate and AZ treatments

$[\text{NaHCO}_3][\text{AZ}]^a$ mmol/L	0	1.0	10.0
1.0	$0.54 \pm 0.06$	$0.39 \pm 0.06$	$0.02 \pm 0.07$
2.5	$0.52 \pm 0.06$	$0.35 \pm 0.06$	$0.04 \pm 0.07$
5.0	$0.31 \pm 0.07$	$0.30 \pm 0.07$	$0.12 \pm 0.07$
20.0	$0.18 \pm 0.07$	$0.14 \pm 0.08$	0

<sup>a</sup> $\text{NaHCO}_3$  or AZ concentration added in the culture medium separately

The  $\delta^{13}\text{C}$  values of the two kinds of  $\text{NaHCO}_3$  added are  $-17.4\%$  or  $-28.4\%$  separately( $\%$ , PDB)

## 4 Discussion

### 4.1 The effect of bicarbonate and acetazolamide on microalgae growth and carbon isotopes

With increasing  $\text{HCO}_3^-$  added to the culture medium, growth of the microalgae increased. It had already been widely confirmed that  $\text{HCO}_3^-$  can stimulate algal growth (Wu et al. 2012; White et al. 2013; Xie and Wu 2017). However, the growth of the treated microalgae decreased sharply with the increase in AZ added to the culture medium since AZ is a CA inhibitor that selectively inhibits CAex activity (Moroney et al. 1985). With AZ, the CAex that catalyzes the reversible interconversion between  $\text{HCO}_3^-$  and  $\text{CO}_2$  was inhibited. Thus, microalgae growth was delayed. The result is that microalgae biomass decreased significantly with addition of AZ.

Simultaneously, the stable carbon isotope composition in algae reflects the utilization of DIC (Chen et al. 2009). In


this study, the more negative the added  $\text{NaH}^{13}\text{CO}_3$ , the more negative the  $\delta^{13}\text{C}$  detected in microalgae for the same concentration of added  $\text{NaHCO}_3$ . Stable carbon isotope composition in the microalgae and  $\text{HCO}_3^-$  concentration in the medium were negatively correlated, and especially so when  $\text{NaH}^{13}\text{CO}_3$  ( $-28.4\%$ ) was added. This demonstrates that the utilization of DIC can alter the  $\delta^{13}\text{C}$  value in microalgae. Moreover, the stable carbon isotope composition without AZ was significantly different from that with AZ. The  $\delta^{13}\text{C}$  value in microalgae with AZ was more negatively altered than that without AZ. The  $\delta^{13}\text{C}$  in microalgae was the most negative in the presence of 10.0 mmol/L AZ. This suggests that CAex can alter the utilization of DIC, and accelerate the rapid interconversion between  $\text{HCO}_3^-$  and  $\text{CO}_2$ , because the slow (uncatalyzed) interconversion of  $\text{CO}_2$  would bring approximately 10‰ stable carbon isotope fractionation (Mook et al. 1974).

### 4.2 The effect of bicarbonate and acetazolamide on microalgae carbon utilization

The proportion of the added  $\text{HCO}_3^-$  increased with the level of additional  $\text{HCO}_3^-$  (Table 4). However, the proportion of the  $\text{HCO}_3^-$  pathway decreased in parallel with the increase in additional  $\text{HCO}_3^-$  (Table 5), as CAex was also inhibited by high concentrations of added  $\text{NaHCO}_3$  (Wu et al. 2012). In addition, the proportion of the  $\text{HCO}_3^-$  pathway decreased with increasing AZ. These results demonstrate that CAex boosted the proportion of the  $\text{HCO}_3^-$  pathway.

The pH of karst lakes in southwest China is generally approximately 8.1.  $\text{CO}_2$  in the water is limited—usually less than 1% of total DIC (Riebesell et al. 1993). Under these conditions, bicarbonate is the main form of DIC (Fig. 1). However, the rate of direct bicarbonate utilization

**Fig. 1** Proposed model of dissolved inorganic carbon utilization by algae in karst lake. Note CA, carbon anhydrase. CAex, the extracellular carbon anhydrase. AE, anion exchange



by anion exchange is slow in microalgae (Fig. 1). Compared with the direct utilization of CO<sub>2</sub> and bicarbonate by microalgae without CAex, the major pathway converting CO<sub>2</sub> from bicarbonate is rapidly catalyzed by CAex (Fig. 1). CAex accelerated photosynthetic Ci assimilation, promoted the conversion of bicarbonate to CO<sub>2</sub> for algal physiological needs, and constantly assimilated CO<sub>2</sub> from the atmosphere into the water. Ultimately, we found that the algae used atmospheric CO<sub>2</sub> as the main DIC source via the bicarbonate pathway under the catalysis of CAex (Fig. 1).

In the natural water of karst areas, microalgae have adjusted their Ci metabolism strategy to adapt to the environment. This study found that microalgal utilization of the bicarbonate that preexisted in the water at the karst area was very small whether AZ was added or not. In Lake Hongfeng, the dominant family is Chlorophyceae, which has high CA activity (Wu et al. 2008). When AZ was added in the medium, CAex activity and growth of the dominant microalgae species were inhibited. As 2.5 mmol/L NaHCO<sub>3</sub> was added—which is similar to natural conditions in karst areas—both growth and carbon sequestration capacity of the microalgae were largely suppressed by 10.0 mmol/L AZ (37% compared to that without AZ). This shows that microalgae with CAex have greater carbon sequestration capacity in karst lakes.

## 5 Conclusion

The bidirectional labeling method presented in this study is an effective way to quantify the proportions of Ci sources and their utilization pathways in microalgae. It can help delineate the mechanism of Ci utilization in microalgae under different conditions. In the natural water of karst areas, microalgae used less bicarbonate preexisting in the aqueous medium than CO<sub>2</sub> derived from the atmosphere. CAex generally increased the utilization of Ci from the atmosphere. The microalgae with CAex had greater carbon sequestration capacity in the lake.

**Acknowledgements** This work was supported by the National Natural Sciences Foundation of China (U1612441), Foundation of Guizhou Province ([2014] 2131) and Doctor Foundation of Guizhou Normal University (0514014).

## References

- Axelsson L, Ryberg H, Beer S (1995) Two modes of bicarbonate utilization in the marine green macroalga *Ulva lactuca*. *Plant Cell Environ* 18:439–445
- Bade DL, Pace ML, Cole JJ, Carpenter SR (2006) Can algal photosynthetic inorganic carbon isotope fractionation be predicted in lakes using existing models? *Aquat Sci* 68:142–153
- Badger MR, Price GD (1994) The role of carbonic anhydrase in photosynthesis. *Annu Rev Plant Biol* 45:369–392
- Chen Z, Cheng HM, Chen XW (2009) Effect of Cl<sup>-</sup> on photosynthetic bicarbonate uptake in two cyanobacteria *Microcystis aeruginosa* and *Synechocystis* PCC6803. *Chin Sci Bull* 54:1197–1203
- Colman B, Huertas IE, Bhatti S, Dason JS (2002) The diversity of inorganic carbon acquisition mechanisms in eukaryotic microalgae. *Funct Plant Biol* 29:261–270
- Elzenga JTM, Prins HBA, Stefels J (2000) The role of extracellular carbonic anhydrase activity in inorganic carbon utilization of *Phaeocystis globosa* (Prymnesiophyceae): A comparison with other marine algae using the isotopic disequilibrium technique. *Limnol Oceanogr* 45:372–380
- Fry B, Sherr EB (1984) δ<sup>13</sup>C measurements as indicators of carbon flow on marine and freshwater ecosystems. *Contrib Mar Sci* 27:13–47
- Giordano M, Beardall J, Raven JA (2005) CO<sub>2</sub> concentrating mechanisms in algae: Mechanisms, environmental modulation, and evolution. *Annu Rev Plant Biol* 56:99–131
- Marlier JF, O’Leary MH (1984) Carbon kinetic isotope effects on the hydration of carbon dioxide and the dehydration of bicarbonate ion. *J Am Chem Soc* 106:5054–5057
- Moazami-Goudarzi M, Colman B (2011) Inorganic carbon acquisition in two green marine *Stichococcus* species. *Plant Cell Environ* 34:1465–1472
- Mondal M, Khanra S, Tiwari ON, Gayen K, Halder GN (2016) Role of carbonic anhydrase on the way to biological carbon capture through microalgae—a mini review. *Environ Progress Sustain Energy* 35(6):1605–1615
- Mook WG, Bommerson JC, Staverman WH (1974) Carbon isotope fractionation between dissolved bicarbonate and gaseous carbon dioxide. *Earth Planet Sci Lett* 22(2):169–176
- Moroney JV, Husic HD, Tolbert N (1985) Effect of carbonic anhydrase inhibitors on inorganic carbon accumulation by *Chlamydomonas reinhardtii*. *Plant Physiol* 79:177–183
- Moulin P, Andriá JR, Axelsson L, Mercado JM (2011) Different mechanisms of inorganic carbon acquisition in red macroalgae (Rhodophyta) revealed by the use of TRIS buffer. *Aquat Bot* 95:31–38
- Riebesell U, Wolf-Gladrow DA, Smetacek V (1993) Carbon dioxide limitation of marine phytoplankton growth rates. *Nature* 361:249–251
- Sedmak JJ, Grossberg SE (1977) A rapid, sensitive, and versatile assay for protein using Coomassie brilliant blue G250. *Anal Biochem* 79:544–552
- Smith-Harding TJ, Beardall J, Mitchell JG (2017) The role of external carbonic anhydrase in photosynthesis during growth of the marine diatom *Chaetoceros muelleri*. *J Phycol* 53(6):1159–1170
- Sültemeyer D (1998) Carbonic anhydrase in eukaryotic algae: characterization, regulation, and possible function during photosynthesis. *Can J Bot* 76:962–972
- Talling JF (1976) The depletion of carbon dioxide from lake water by phytoplankton. *J Ecol* 64(1):79–121
- White DA, Pagarette A, Rooks P, Ali ST (2013) The effect of sodium bicarbonate supplementation on growth and biochemical composition of marine microalgae cultures. *J Appl Phycol* 25(1):153–165
- Williams TG, Turpin DH (1987) The role of external carbonic anhydrase in inorganic carbon acquisition by *Chlamydomonas reinhardtii* at alkaline pH. *Plant Physiol* 83:92–96
- Wu YY, Li PP, Wang BL, Liu CQ, He M, Chen CH (2008) Composition and activity of external carbonic anhydrase of microalgae from karst lakes in China. *Phycol Res* 56:76–82

- Wu YY, Xu Y, Li HT, Xing DK (2012) Effect of acetazolamide on stable carbon isotope fractionation in *chlamydomonas reinhardtii* and *chlorella vulgaris*. *Chin Sci Bull* 57:786–789
- Xie TX, Wu YY (2017) The biokarst system and its carbon sinks in response to pH changes: a simulation experiment with microalgae. *Geochem Geophys Geosyst* 18:827–843