ORIGINAL ARTICLE

Translocation and distribution of mercury in biomasses from subtropical forest ecosystems: evidence from stable mercury isotopes

Yi Liu^{1,2} · Che-Jen Lin^{1,3,4} · Wei Yuan^{1,2} · Zhiyun Lu⁵ · Xinbin Feng¹

Received: 25 September 2020/Revised: 25 September 2020/Accepted: 15 October 2020/Published online: 28 October 2020 © Science Press and Institute of Geochemistry, CAS and Springer-Verlag GmbH Germany, part of Springer Nature 2020

Abstract To understand its source, distribution, storage, and translocation in the subtropical forest ecosystems, mercury (Hg) concentrations and stable isotopes in forest biomass tissues (foliage, branch, bark, and trunk) were investigated at Ailao Mountain National Nature Reserve, Southwest China. The total Hg (THg) concentrations in the samples show the following trend: mature foliage $(57 \pm 19 \text{ ng g}^{-1}) > \text{bark}$ $(11 \pm 4.0 \text{ ng g}^{-1}) > \text{branch}$ $(5.4 \pm 2.5 \text{ ng g}^{-1}) > \text{trunk} (1.6 \pm 0.7 \text{ ng g}^{-1})$. Using the measured THg concentrations and the quantity of respective biomasses, the Hg pools in the forest are: wood $(60 \pm 26 \ \mu g \ m^{-2}) > bark$ $(51 \pm 18 \ \mu g \ m^{-2}) > foliage$ $(41 \pm 11 \ \mu g \ m^{-2}) > branch \ (26 \pm 8.3 \ \mu g \ m^{-2})$. The tree biomasses displayed negative δ^{202} Hg (-1.83‰ to -3.84%) and Δ^{199} Hg (-0.18% to -0.62%). The observed Δ^{200} Hg (- 0.08% to 0.04%) is not significantly from zero. A Δ^{199} Hg/ Δ^{201} Hg ratio of 1.05 was found in tree biomasses, suggesting that mercury has undergone Hg^(II) photoreduction processes. A Hg-isotope based binary

Che-Jen Lin Jerry.Lin@lamar.edu

Xinbin Feng fengxinbin@vip.skleg.cn

- ¹ State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang, China
- ² University of Chinese Academy of Sciences, Beijing, China
- ³ Center for Advances in Water and Air Quality, Lamar University, Beaumont, TX, USA
- ⁴ College of Environment and Energy, South China University of Technology, Guangzhou, China
- ⁵ National Forest Ecosystem Research Station, Ailaoshan, Yunnan, China

mixing model suggests that Hg in the tree biomasses mainly originated from foliage uptake of atmospheric Hg⁰, constituting 67%, 80%, and 77% of Hg in wood, branch, and bark, respectively. Our study sheds new light on the transportation and sources of Hg in the subtropical forest ecosystems.

Keywords Hg \cdot Subtropical forest ecosystem \cdot Hg stable isotope \cdot Hg-isotope based binary mixing model

1 Introduction

Mercury (Hg) in the atmosphere mainly exists (> 90%) in the form of gaseous elemental Hg (GEM) (Gratz et al. 2010), whose atmospheric lifetime (0.5 - 1 year) allows long-range transport globally (Selin 2009). Forest ecosystems play an important role in the biogeochemical cycling of Hg on a global scale. Oxidation of Hg⁰ followed wet deposition (e.g., precipitation and/or throughfall), or Hg⁰ uptake by vegetation followed by litterfall deposition are the main pathways for Hg deposition. Recent studies suggested that the forest ecosystems act as a major sink of atmospheric Hg⁰ through litterfall Hg deposition (Johnson and Lindberg 1995; Laacouri et al. 2013; Louis et al. 2001; Wang et al. 2017). However, the post-depositional biogeochemical processes controlling the storage, accumulation, and sequestration of Hg remain not well understood (Gu et al. 2011; Jiang et al. 2015; Jiskra et al. 2015; Wang et al. 2003; Yu et al. 2016; Yuan et al. 2019).

The Hg distribution in forest ecosystems highly depends on the processes governing the translocation of Hg after vegetative uptake, while the translocation and distribution result in the storage of Hg in the forest. Previous studies reported the levels of THg in plants (Ericksen et al. 2003; Kang et al. 2019; Yang et al. 2018), suggesting that a dual mechanism controlled Hg uptake in tree tissues with some uptake from the soil by the roots and the majority of uptake from atmospheric sources by the leaves. A recent study reported that the Hg pool size in tree woody biomass can be 1-3 times higher than that in foliage at four forested sites in the northeastern USA, which highlighted the role of woody biomass in Hg cycling in forest ecosystems (Yang et al. 2018).

Hg isotopic signatures provide a valuable fingerprint for tracing source and process (Bergquist and Blum 2009; Blum et al. 2014; Yin et al. 2014). The isotopic composition of Hg in foliage in the forest ecosystem has been investigated extensively (Demers et al. 2013a; Yu et al. 2016; Yuan et al. 2019; Zheng et al. 2016), explaining Hg in the foliage was mainly sourced from GEM. However, data suggesting the sources and net reserves of Hg in biomass tissues of trees other than foliage are still relatively rare, which limits the understanding of Hg cycling in forest ecosystems.

The objective of this study is to provide a comprehensive description of the distribution, storage, and translocation of Hg in biomass in a subtropical forest ecosystem, the Ailao Mountain National Nature Reserve, Southwest China. We performed measurements on THg concentrations, and its associated isotopic compositions in foliage, branch, bark, and trunk biomasses and estimate the Hg storage pools in each of the biomass compartments. These measurements aim to gain a better understanding of the distribution and sources of Hg in the forest ecosystem.

2 Materials and methods

2.1 Study area, sampling and sample preparation

The Xujiaba region $(24^{\circ}32'N, 101^{\circ}01'E)$ is located in the Ailao Mountain National Nature Reserve, Southwest China, at an altitude of 2400–2650 m (You 1983) (Fig. 1). It is a subtropical evergreen broad-leaved forest characterized with abundant rainfall (1840 mm per year) (Song et al. 2015) with three dominant evergreen tree species, including *Lithocarpus xylocarpus (LX)*, *Castanopsis wattii (CW)*, and *Schima Noronhae (SN)* at the proportion of 16.3%, 16.3%, and 23.3%, respectively (Xie et al. 1996). The frost-free period lasts about 180 days annually (Tan et al. 2011). Influenced by the Indian Ocean southwest monsoon, the forest is cool and humid, with an annual average temperature and relative humidity at 11.3 °C and 84%, respectively. In the forest area, the soil is primarily yellow–brown with a pH of 4.4–4.9 (Liu et al. 2002).

Samples were collected at three elevations (S₁: N24.5398°, E101.0278°, 2450 m a.s.l; S₂: N24.53817°,

E101.02968°, 2550 m a.s.l; S₃: N24.53629°, E102.03129°, 2650 m a.s.l). Three dominant tree species as previously mentioned were selected for biomass sampling. Mature foliage, branch, bark, and wood were sampled in October 2016. The foliage and branch samples were sampled at three canopy heights (i.e., upper, middle, and lower canopy) at three heights (20, 21, and 22 m), and then mixed as a composite sample. The bark and wood samples from each selected tree species were collected at the chest height (~ 1.5 m) at three replicates. Bedrock samples were sampled from a depth of 20 m underground using a drilling machine.

All the samples were placed in clean polyethylene bags and taken back to the laboratory for analysis. During the each sampling, new polyethylene gloves were worn to avoid cross-contamination. Each sample was washed by DDW and freeze-dried at -60° to constant weight, then ground to fine powder.

2.2 Sample analysis

2.2.1 THg concentration

THg concentration in tree biomass samples was measured by the Lumex RA-915+ (Lumex Analytics, Russia, detection limit: 1 ng g⁻¹) atomic absorption spectrometric analyzer (Wang et al. 2010). The method recovery was $4.6 \pm 3.5 \%$ (n = 12, 1 σ), 96.3 $\pm 5.4 \%$ (n = 10, 1 σ), 97.2 $\pm 4.6 \%$ (n = 10, 1 σ) for GBW10049 (GSB-27), BCR-482 (Lichen) and GBW10020 (GSB-11) respectively.

2.2.2 Hg isotopes

The samples were prepared by pre-concentration following a previous method (Sun et al. 2013). Briefly, Hg was preconcentrated by double-stage tube furnace combustion, and the vaporized Hg⁰ was subsequently trapped by a 5 mL freshly 40 % concentrated mixture of HNO3 and HCl (HNO₃: HCl = 2:1, V/V). Before Hg isotope measurement, THg concentration of all trapping solutions was analyzed by cold vapor atomic fluorescence spectrometry (CVAFS, Tekran 2500, Canada) using the US-EPA method 1631 (1631, 2002). The recovery of Hg was $97.2 \pm 6.3 \%$ $(n = 5, mean \pm 1\sigma)$ during thermolysis the standard reference material BCR482 (lichen, $480 \pm 20 \text{ ng g}^{-1}$), and the recoveries of Hg were $92.5 \pm 4.3\%$ (n = 9, mean \pm 1σ) for foliage samples, 90.6 \pm 3.2 % (n = 6, mean \pm 1 σ) for branch samples, 93.7 \pm 2.3 % (n =4, mean \pm 1 σ) for bark samples, $91.8 \pm 3.5 \%$ (n = 4, mean $\pm 1\sigma$) for wood samples, and 90.7 \pm 3.2 % (n = 4, mean \pm 1 σ) for rock samples respectively, according to THg measured by Lumex RA-915 +. Hg isotopic compositions were determined by a Nu Plasma II Multiple-collector inductively

Fig. 1 Site locations for biomasses collection at different altitudes from Mt. Ailao. S_1 , S_2 and S_3 represent three different elevations in forest respectively (S1: N24.5398°, E101.0278°, 2450 m a.s.l; S2: N24.53817°, E101.02968°, 2550 m a.s.l; S3: N24.53629°, E102.03129°, 2650 m a.s.l)



coupled plasma mass spectrometry (Nu Instruments, UK), following a previous method (Yin et al. 2010).

Mass dependent fractionation (MDF) and mass independent fractionation (MIF) of Hg isotopes, in units of permil (‰), were expressed as the following equations (Bergquist and Blum 2007b):

$$\delta^{xxx}Hg(_{00}) = \left\{ \left[\left({^{xxx}Hg/^{198}Hg} \right)_{sample} / \left({^{xxx}Hg/^{198}Hg} \right)_{NIST-3133} \right] - 1 \right\} \times 1000$$

$$(1)$$

$$\Delta^{\text{xxx}}\text{Hg} = \delta^{\text{xxx}}\text{Hg} - \beta^{\text{xxx}} \times \delta^{202}\text{Hg}$$
(2)

xxx is the mass of each Hg isotope from 199 to 202 amu, the factor β^{xxx} is 0.252 for ¹⁹⁹Hg, 0.502 for ²⁰⁰Hg and 0.752 for ²⁰¹Hg, respectively.

During the Hg isotopes analysis, in addition to repetitive measurement (typically n = 2), we also used the UM-Almadén as a second standard and measured once in every 10 samples. Results of UM-Almadén ($-0.52 \pm 0.06 \%$ for δ^{202} Hg, $0.03 \pm 0.05 \%$ for Δ^{199} Hg, $-0.01 \pm 0.05 \%$ for Δ^{200} Hg and $-0.02 \pm 0.10 \%$ for Δ^{201} Hg, 2SD, n = 5) and BCR-482 ($-1.53 \pm 0.22 \%$ for δ^{202} Hg, -0.66 ± 0.08 for Δ^{199} Hg, $-0.06 \pm 0.03 \%$ for Δ^{200} Hg and $-0.67 \pm 0.06 \%$ for Δ^{201} Hg, 2SD, n = 5) were consistent with the recommended values (Bergquist and Blum 2007a; Blum and Bergquist 2007; Estrade et al. 2010).

3 Results and discussion

3.1 Distribution of THg in biomass

The measured THg concentrations (Fig. 2) in matured foliage ranged from 29.6 to 105.9 ng g⁻¹. THg concentrations in the fresh litter were 74 ± 19 ng g⁻¹ for CW (mean \pm SD, n = 9), 56 \pm 13 ng g⁻¹ for LX (mean \pm



Fig. 2 THg concentrations measured in the vegetative samples in this study

SD, n = 9), and 42 ± 10 ng g⁻¹ for SN (mean ± SD, n = 9), respectively. The varied foliage THg was attributed to tree species that yield different uptake flux and reemission flux (Du et al. 2018; Louis et al. 2001; Luo et al. 2016; Poissant et al.2008; Yuan et al. 2019). The assimilated Hg in foliage is expected to be transferred to the trunk (Frescholtz et al. 2003). Hg^{II} in the soil can be utilized and moved through the root (Ericksen et al. 2003; Rea et al. 2002), which only play the secondary role in accumulating Hg in-ground biomass (Cui et al. 2014). The observed THg concentrations were 5.4 ± 2.5 ng g⁻¹ for branch (mean ± SD, n = 27), 11 ± 4.0 ng g¹ for bark (mean ± SD, n = 27).

Table 1 shows THg concentration of plant tissues in forest ecosystems observed at global sites. Previous studies (JA et al. 2003; Kang et al. 2019; Siwik et al. 2009; Sun et al. 2017; Yang et al. 2018; Zhou et al. 2016, 2017) have focused on the distribution of Hg in foliage and trunk. The data show a wide range of THg concentrations in foliage from 3 to 160 ng g^{-1} . The large variation may be the resulted from the tree species, which were verified in previous studies (Juillerat et al. 2012; Richardson and Friedland 2015; Siwik et al. 2009); foliage age, which resulted in long exposure and accumulation time in longevity foliage (Ericksen et al. 2003; JA et al. 2003; Millhollen et al. 2006; Rea et al. 2002; Selvendiran et al. 2008; Siwik et al. 2009); climate conditions and atmospheric THg concentrations, which affected the THg accumulation in foliage (Frescholtz et al. 2003). These factors influence

uptake and absorption by stoma (Demers et al. 2013b; Laacouri et al. 2013; Yuan et al. 2019; Zheng et al. 2016). The tree trunk has low levels of THg concentration (Table 1). In most areas, the THg levels in the trunk were below 5 ng g⁻¹ (Kang et al. 2019; Siwik et al. 2009; Yang et al. 2017, 2018). Our results indicate that THg content in the trunks of the subtropical evergreen broad-leaved forest from Ailao Mountain ranged from 0.8 to 2.4 ng g⁻¹. Data for bark and branch, constituting 7 %–10 % of biomasses in forest ecosystems, are rarely reported. Available data suggested bark contained significantly higher THg than branch (Zhou et al. 2016, 2017). Our study shows a similar trend (Fig. 2), suggesting that despite differences in sampling heights and tree species, the mechanism of Hg retention in tree branch and bark was similar.

Globally, forest biomass and soil constitute some of the largest reservoirs of actively cycling Hg (Juillerat et al. 2012), especially in tropical and subtropical forest ecosystems (Wang et al. 2019). Atmospheric Hg⁰ could be taken up by foliage via stomata or absorbed on the surface (Laacouri et al. 2013; Yuan et al. 2019). Further, atmospheric Hg was assimilated by foliage and partly released ($\sim 30\%$) again to the atmosphere (Yuan et al. 2019), resulting in a large proportion of Hg accumulation. The accumulated Hg in foliage could be transferred to the trunk based in a previous study (Frescholtz et al. 2003), while another study showed the opposite results that THg concentrations in branch and trunk were influenced by THg level in soil (Ericksen et al. 2003). Meanwhile, the Hg^{II} in the soil can be utilized and moved through the root

Location	Foliage	Branch	Bark	Wood	References
China					
AiLao Mt	29.6-105.9	5.4 ± 2.5	11 ± 4.0	1.6 ± 0.7	This study
Tianshan	11.8-20.7		14.0–19.4	3.0-4.6	Kang et al. (2019)
Tieshanping	74–210	12–23	22-34	8–28	Zhou et al. (2016)
Dongling Mt	26.0-65.9	7.1–25.3	2.7-35.6	0.6–7.3	Zhou et al. (2017)
Jiulong Estuary	17.7–28.4			4.0-10.6	Sun et al. (2017)
New Hampshire				1.48-4.59	Yang et al. (2017)
Northeastern	16.3-28.6		7.7–22.5	1.4-2.8	Yang et al. (2018)
USA					
Northeast				8 ± 4	Richardson and Friedland (2015)
Huntington	4.4-37.3				Bushey et al. (2008)
Reno NV	30-160	~ 20		~ 20	Ericksen et al. (2003)
Guiana	59–70				Mélières et al. (2003)
French					
Amiata Mt			5–98		Chiarantini et al. (2016)
Italy					
Ontario	3.0-45.3			2	Siwik et al. (2009)
Canada					
Saskatcheman	8		16–38		Friedli et al. (2007)

Table 1	THg concentration of
biomass	in related study sites



Fig. 3 Isotopic signature of Hg for the vegetative and bed rocks in this study

(Ericksen et al. 2003). Such a process that occurred at root is expected to be a secondary process in accumulating Hg in-ground biomass such as branch and trunk.

3.2 Storage of THg in biomass

Given the large biomass mass in bark, branch, and trunk in a forest with a low turnover rate due to a relatively long life cycle, the storage and transformation of Hg in such biomass could play a significant role in Hg cycling and should not be ignored. At Mt. Ailao, the average ground biomass was 7.1 t ha⁻¹, 33 t ha⁻¹, 46 t ha⁻¹, 373 t ha⁻¹ for foliage, branch, bark, and trunk, respectively. The pools size of Hg was estimated using the following equation:

$$\text{Pool}_{\text{biomass}} = C_{\text{b}} \times M_{\text{b}}.$$
(3)

where, C_b (ng g⁻¹) and M_b (t ha⁻¹) are the THg concentration and biomass of tree biomass samples, respectively (Zhou et al. 2017). Table 2 shows the average pools of Hg in biomass followed in a descending order: wood (60 µg m⁻²) > bark (51 µg m⁻²) > foliage (41 µg m⁻²) > branch (26 µg m⁻²). The corresponding proportion was 33.7 %, 28.7 %, 23.0 %, and 14.6 % for the pool of biomass in wood, bark, foliage, and branch, separately. This distribution is different from the reported biomass pool (Zhou et al. 2017) reported previously, which a pool size in

the order of branch > foliage > bole > bark for Chinese pine, branch > bole > bark > foliage for a mixed broadleaf forest. For Larch and Oak, the order was similar: branch > bark > foliage > bole. These suggested that the tissue of the branch was the largest pool of Hg; and that the Hg reservoirs vary in different forest types due to the biomass affected by the canopy geometry, areal density of species, and individual differences of each plant (Bushey et al. 2008; Richardson and Friedland 2015) and the THg concentration probably due to the physiological property of plants, such as migration rate, and retention in the xvlem (Zhou et al. 2017). Although wood has the lowest THg concentration at $< 2 \text{ ng g}^{-1}$, it has the largest pool size of Hg due to its greatest biomass at 60 μ g m⁻². The bark is the second large Hg reservoir due to the higher THg content and greater biomasses, up to 51 μ g m⁻². The foliage has a slightly lower pool of Hg than bark at 41 μ g m⁻², while the branch has the smallest pool of Hg at $26 \ \mu g \ m^{-2}$. The ground biomass of the subtropical forest ecosystem store a large quantity of Hg mass at nearly 180 $\mu g m^{-2}$.

3.3 Sources and processes of Hg implicated from stable isotope data

Characteristics of Hg stable isotope in the forest ecosystem provided a useful fingerprint for tracing Hg translocation and transformation. As listed in Table 3, all the biomass samples exhibited distinctive negative δ^{202} Hg (- 1.83 to -3.84 ‰) and Δ^{199} Hg (-0.18 ‰ to -0.62 ‰), but near-zero Δ^{200} Hg (- 0.08 ‰ to 0.04 ‰). For foliage, Hg isotopic composition was characterized with negative δ^{202} Hg and Δ^{199} Hg averaged with -2.95 ± 0.96 ‰ and -0.35 ± 0.13 %, respectively, while neglectable Δ^{200} Hg averaged with $-0.03 \pm 0.05 \%$ (n = 9, 2 σ). Such results were consistent with Hg isotopic compositions in leaves reported earlier (Enrico et al. 2016; Fu et al. 2016; Yu et al. 2016; Yuan et al. 2019), which was attributed to the uptake process that causes preferential absorption of lighter isotopes. Subsequently, the reemission process that occurred at leaves produces negative odd-MIF remained in the residue (Yuan et al. 2019).

Hg isotopic compositions in-branch averaged - 2.80 \pm 0.90 % and - 0.35 \pm 0.21 % for $\delta^{202} Hg$ and

Sample type	n Biomass/(t ha ⁻¹)		THg/(ng g	g ⁻¹)	Hg pool siz	Hg pool size/($\mu g m^{-2}$)	
			Mean	SD	Mean	SD	
Foliage	27	7.1	57	16	41	11	
Branch	27	33	5.4	2.5	26	8.3	
Bark	27	46	11	4.0	51	18	
Wood	27	373	1.6	0.7	60	26	

 Table 2
 THg concentration in vegetative biomasses and their associated storage

 Δ^{199} Hg Δ^{200} Hg respectively. while averaged 0.01 ± 0.09 ‰ (n = 6, 2 σ). Bark showed negative δ^{202} Hg and Δ^{199} Hg as -2.46 ± 1.13 ‰ and -0.44 ± 0.19 ‰ respectively, but negligible Δ^{200} Hg as -0.05 ± 0.07 ‰ $(n = 4, 2\sigma)$. Wood also displayed negative δ^{202} Hg and Δ^{199} Hg with -2.72 ± 0.46 ‰ and -0.35 ± 0.12 ‰ Δ^{200} Hg independently, but insignificant with -0.01 ± 0.04 ‰ (n = 4, 2 σ). The similar MIF values among foliage, branch, bark, and trunk (Fig. 3) revealed Hg in biomasses were mainly derived from foliage transport, and there was no MIF generated during transmission (Sun et al. 2017; Yin et al. 2013a). The slope of Δ^{199} Hg/ Δ^{201} Hg is a useful fingerprint to identify the geochemical processes since the transformation in the tree does not shift Hg odd-MIF (Estrade et al. 2009; Ghosh et al. 2013; Jiskra et al. 2012; Koster van Groos et al. 2014; Wiederhold et al. 2010; Yin et al. 2013b). The slope of Δ^{199} Hg/ Δ^{201} Hg in all the samples is found to be $1.05 (R^2 = 0.86)$, suggesting that mercury has undergone Hg(II) photoreduction before entering the tree biomass (Fig. 4).

Given the Hg in-ground biomass were mainly contributed by atmospheric Hg⁰ uptake through foliage and bedrock through the root (Sun et al. 2017; Yin et al. 2013a), the contribution of foliage and bedrock can be calculated. Different from the MDF of Hg isotopes, which can be caused in almost all processes, including chemical processes (Bergquist and Blum 2007a; Sun et al. 2016; Yang and Sturgeon 2009; Zheng et al. 2019; Zheng and Hintelmann 2009, 2010b), physical process (Estrade et al. 2009; Ghosh et al. 2013; Jiskra et al. 2012; Koster van Groos et al. 2014; Wiederhold et al. 2010; Yin et al. 2013b), and biogeochemical process in which microorganisms were involved (Bergquist and Blum 2007a; Kritee et al. 2008, 2007, 2013; Malinovsky and Vanhaecke 2011; Perrot et al. 2015), the odd-MIF of Hg isotopes is



Fig. 4 $\bigtriangleup^{199} \rm Hg$ versus $\bigtriangleup^{201} \rm Hg$ for the vegetative samples in this study

considered to be caused only in some special geochemical processes, such as photochemical process, methylmercury photodegradation, equilibrium evaporation and organic matters dark reduction/ oxidation (Bergquist and Blum 2007a; Blum et al. 2013; Estrade et al. 2009; Ghosh et al. 2013; Jimenez-Moreno et al. 2013; Jiskra et al. 2012; Wiederhold et al. 2010; Yang and Sturgeon 2009; Zheng et al. 2019; Zheng and Hintelmann 2009, 2010a, 2010b). Therefore, odd-MIF is utilized to build a two end-member mixing model to quantify of the source of Hg in-ground biomass in the following equation based on a previous study (Yin et al. 2013a).

$$\Delta^{199} \text{Hg}_{\text{biomass}} = F_{\text{foliage}} \Delta^{199} \text{Hg}_{\text{f}} + (1 - F_{\text{foliage}}) \Delta^{199} \text{Hg}_{\text{rock}}$$
(4)

$$F_{\text{foliage}}(\%) = \left(\Delta^{199} \text{Hg}_{\text{biomass}} - \Delta^{199} \text{Hg}_{\text{rock}}\right) \\ \times 100 / \left(\Delta^{199} \text{Hg}_{\text{foliage}} - \Delta^{199} \text{Hg}_{\text{rock}}\right)$$
(5)

where Δ^{199} Hg_{biomass} is the Δ^{199} Hg values of biomass including samples of the branch, bark, and wood. Δ^{199} Hg foliage and Δ^{199} Hg_{rock} are the Δ^{199} Hg values of foliage Hg and rock Hg, respectively. F_{foliage} (%) is the fraction of foliage Hg source. Considering the uncertainties that occurred at isotopic measurement, to further assess the uncertainty in calculation results, a Monte Carlo simulation was applied in the modeling calculation. More specifically, the every Hg isotopic signature, which was used in solving the above equations, was produced randomly around the mean with 1 standard error followed a normal distribution. Subsequently, the equation would be solved to get the fraction of different end-members. Such operations were applied to one million simulations to accomplish the simulation of a fraction of foliage or bedrock and its uncertainties. Results from the end-member mixing model (Fig. 5) showed that atmospheric Hg followed by foliage



Fig. 5 Estimated fraction of atmospheric Hg in the vegetative biomass samples

Table3 Isotop	ic comp	osition of m	ercury me	easured in gr	ound tree	biomasses	and rocks									
Samples	u	$\delta^{199} Hg$	2sd	δ^{200} Hg	2sd	δ^{201} Hg	2sd	δ^{202} Hg	2sd	${\rm g}{\rm H}^{99}{\rm H}{\rm g}$	2sd	$\Delta^{200}{ m Hg}$	2sd	$\Delta^{201} \mathrm{Hg}$	2sd	Average TH _α
		%00	%00	%00	%00	%o	%00	%00	%00	%00	%00	%00	%00	%00	%00	ng g ⁻¹
Foliage	6	- 1.09	0.33	- 1.51	0.46	- 2.56	0.79	- 2.95	0.96	- 0.35	0.13	-0.03	0.05	- 0.34	0.14	48
Branch	9	-1.06	0.23	-1.40	0.39	- 2.44	0.56	-2.80	06.0	-0.35	0.21	0.01	0.09	-0.34	0.19	5.4
Bark	4	-1.06	0.39	- 1.29	0.47	- 2.24	0.91	- 2.46	1.13	- 0.44	0.29	-0.05	0.07	-0.39	0.24	11
Wood	4	- 1.03	0.17	- 1.38	0.26	- 2.37	0.28	-2.72	0.46	-0.35	0.12	-0.01	0.04	-0.33	0.19	1.6
Rock	4	-0.17	0.07	-0.14	0.09	-0.32	0.17	-0.29	0.24	-0.10	0.09	0.01	0.04	-0.10	0.03	5.2
-MU	5	-0.16	0.06	-0.25	0.08	-0.41	0.18	-0.52	0.06	-0.03	0.05	-0.01	0.05	- 0.02	0.10	1.0
Almadén																
BCR-482	5	-1.04	0.11	-0.72	0.12	- 1.81	0.20	- 1.53	0.22	-0.66	0.08	-0.06	0.03	-0.67	0.06	467

transport was the dominant source for the ground woody biomass. Approximately 66.7 % of the Hg in the wood was derived from Hg transport through the foliage, while Hg in bark and branch were derived mainly from the foliage transport, up to 77 % and 80 % respectively. The much higher atmospheric Hg fractions in the woody biomass suggested that soil contributes little to the ground biomass Hg (Table 3).

4 Conclusion

This paper explores the THg distribution, storage, and Hg isotope composition in each part of over-ground tree biomass in the subtropical evergreen broad-leave forest of Ailao Mountain National Nature Reserve. We concluded that the THg in the biomass samples is derived mainly from the foliage transmission to ground biomass reaching 80 %, 77 %, 66.7 % in branch, bark, and trunk, respectively. The net storage reservoir of Hg approximately is nearly 180 μ g m⁻² in-ground biomasses, suggesting that tree biomass acts as a sink in the Hg cycling in forest ecosystems. Approximately 100 μ g m⁻² was derived from foliage transport to branch, bark, and trunk, making up 80 % of the Hg reservoir in-ground biomasses, while bedrock only contributed 20 % Hg to the ground biomass. These indicate that the transmission of foliage Hg to branch, bark, and trunk is the main route of Hg biogeochemical cycling in forest ecosystems.

Acknowledgement This work was funded by the National Natural Science Foundation of China (No. 41430754). We thank Jin-Hua Qi for the assistance in sample collection. The Ailao Mountain Ecosystem Research Station was thanked for its fieldwork support.

References

- 1631 USEPAM (2002) Revision E: mercury in water by oxidation, purge and trap, and cold vapor atomic fluorescence spectrometry. https://www.epa.gov/sites/production/files/2015-08/docu ments/method_1631e_2002.pdf
- Bergquist BA, Blum JD (2007a) Mass-dependent and -independent fractionation of Hg isotopes by photoreduction in aquatic systems. Science 318:417–420
- Bergquist BA, Blum JD (2007b) Mass-dependent and mass-independent fractionation of Hg isotopes by photo-reduction in aquatic systems. Science 318:417–420
- Bergquist BA, Blum JD (2009) The odds and evens of mercury isotopes: applications of mass-dependent and mass-independent isotope fractionation. Elements 5:353–357
- Blum JD, Bergquist BA (2007) Reporting of variations in the natural isotopic composition of mercury. Anal Bioanal Chem 388:353–359
- Blum JD, Popp BN, Drazen JC, Choy CA, Johnson MW (2013) Methylmercury production below the mixed layer in the North Pacific Ocean. Nat Geosci 6:879–884

- Blum JD, Sherman LS, Johnson MW (2014) Mercury isotopes in earth and environmental sciences. Annu Rev Earth Planet Sci 42:249–269
- Bushey JT, Nallana AG, Montesdeoca MR, Driscoll CT (2008) Mercury dynamics of a northern hardwood canopy. Atmos Environ 42:6905–6914
- Chiarantini L, Rimondi V, Benvenuti M, Beutel MW, Costagliola P, Gonnelli C et al (2016) Black pine (*Pinus nigra*) barks as biomonitors of airborne mercury pollution. Sci Total Environ 569–570:105–113
- Cui L, Feng X, Lin CJ, Wang X, Meng B (2014) Accumulation and translocation of 198Hg in four crop species. Environ Toxicol Chem 33:334–340
- Demers JD, Blum JD, Zak DR (2013a) Mercury isotopes in a forested ecosystem: implications for air-surface exchange dynamics and the global mercury cycle. Glob Biogeochem Cycles 27:222–238
- Demers JD, Blum JD, Zak DR (2013b) Mercury isotopes in a forested ecosystem: implications for air-surface exchange dynamics and the global mercury cycle. Glob Biogeochem Cycles 27:222–238
- Du H, Ma M, Sun T, An S, Igarashi Y, Wang D (2018) Methyl and total mercury in different media and associated fluxes in a watershed Forest, Southwest China. Int J Environ Res Public Health 15:2618–2631
- Enrico M, Le Roux G, Marusczak N, Heimburger LE, Claustres A, Fu XW et al (2016) Atmospheric mercury transfer to peat bogs dominated by gaseous elemental mercury dry deposition. Environ Sci Technol 50:2405–2412
- Ericksen JA, Gustin MS, Schorran DE, Johnson DW, Lindberg SE, Coleman JS (2003) Accumulation of atmospheric mercury in forest foliage. Atmos Environ 37:1613–1622
- Estrade N, Carignan J, Sonke JE, Donard OFX (2009) Mercury isotope fractionation during liquid-vapor evaporation experiments. Geochim Cosmochim Acta 73:2693–2711
- Estrade N, Carignan J, Sonke JE, Donard OFX (2010) Measuring Hg isotopes in bio-geo-environmental reference materials. Geostand Geoanal Res 34:79–93
- Frescholtz TF, Gustin MS, Schorran DE, Fernandez GCJ (2003) Assessing the source of mercury in foliar tissue of quaking aspen. Environ Toxicol Chem 22:2114–2119
- Friedli HR, Radke LF, Payne NJ, McRae DJ, Lynham TJ, Blake TW (2007) Mercury in vegetation and organic soil at an upland boreal forest site in Prince Albert National Park, Saskatchewan, Canada. J Geophys Res Biogeosci 112(G1):G01004. https://doi. org/10.1029/2005JG000061
- Fu XW, Zhu W, Zhang H, Sommar J, Yu B, Yang X, et al. Depletion of atmospheric gaseous elemental mercury by plant uptake at Mt. Changbai, Northeast China. Atmospheric Chemistry and Physics 2016; 16: 12861–12873.
- Ghosh S, Schauble EA, Couloume GL, Blum JD, Bergquist BA (2013) Estimation of nuclear volume dependent fractionation of mercury isotopes in equilibrium liquid-vapor evaporation experiments. Chem Geol 336:5–12
- Gratz LE, Keeler GJ, Blum JD, Sherman LS (2010) Isotopic composition and fractionation of mercury in great lakes precipitation and ambient air. Environ Sci Technol 44:7764–7770
- Gu B, Bian Y, Miller CL, Dong W, Jiang X, Liang L (2011) Mercury reduction and complexation by natural organic matter in anoxic environments. Proc Natl Acad Sci 108:1479–1483
- Ja E, Ms G, De S (2003) Accumulation of atmospheric mercury in forest Foliage. Atmos Environ 37:1613–1622
- Jiang T, Skyllberg U, Wei S, Wang D, Lu S, Jiang Z et al (2015) Modeling of the structure-specific kinetics of abiotic, dark reduction of Hg(II) complexed by O/N and S functional groups in humic acids while accounting for time-dependent structural rearrangement. Geochim Cosmochim Acta 154:151–167

49

- (2013) Chemical kinetic isotope fractionation of mercury during abiotic methylation of Hg(II) by methylcobalamin in aqueous chloride media. Chem Geol 336:26–36
- Jiskra M, Wiederhold JG, Bourdon B, Kretzschmar R (2012) Solution speciation controls mercury isotope fractionation of Hg(II) sorption to goethite. Environ Sci Technol 46:6654–6662
- Jiskra M, Wiederhold JG, Skyllberg U, Kronberg R-M, Hajdas I, Kretzschmar R (2015) Mercury deposition and re-emission pathways in boreal forest soils investigated with hg isotope signatures. Environ Sci Technol 49:7188–7196
- Johnson DW, Lindberg SE (1995) The biogeochemical cycling of Hg in forests: alternative methods for quantifying total deposition and soil emission. Water Air Soil Pollut 80:1069–1077
- Juillerat JI, Ross DS, Bank MS (2012) Mercury in litterfall and upper soil horizons in forested ecosystems in Vermont, USA. Environ Toxicol Chem 31:1720–1729
- Kang H, Liu X, Guo J, Wang B, Xu G, Wu G et al (2019) Characterization of mercury concentration from soils to needle and tree rings of Schrenk spruce (*Picea schrenkiana*) of the middle Tianshan Mountains, northwestern China. Ecol Ind 104:24–31
- Koster van Groos PG, Esser BK, Williams RW, Hunt JR (2014) Isotope effect of mercury diffusion in air. Environ Sci Technol 48:227–233
- Kritee K, Blum JD, Johnson MW, Bergquist BA, Barkay T (2007) Mercury stable isotope fractionation during reduction of Hg(II) to Hg(0) by mercury resistant microorganisms. Environ Sci Technol 41:1889–1895
- Kritee K, Blum JD, Barkay T (2008) Mercury stable isotope fractionation during reduction of Hg(II) by different microbial pathways. Environ Sci Technol 42:9171–9177
- Kritee K, Blum JD, Reinfelder JR, Barkay T (2013) Microbial stable isotope fractionation of mercury: a synthesis of present understanding and future directions. Chem Geol 336:13–25
- Laacouri A, Nater EA, Kolka RK (2013) Distribution and uptake dynamics of mercury in leaves of common deciduous tree species in Minnesota, USA. Environ Sci Technol 47:10462–10470
- Liu W, Fox JED, Xu Z (2002) Biomass and nutrient accumulation in montane evergreen China. For Ecol Manag 158:223–235
- Louis VLS, Rudd JWM, Kelly CA, Hall BD, Rolfhus KR, Scott KJ et al (2001) Importance of the forest canopy to fluxes of methyl mercury and total mercury to boreal ecosystems. Environ Sci Technol 35:3089–3098
- Luo Y, Duan L, Driscoll CT, Xu G, Shao M, Taylor M, et al (2016) Foliage/atmosphere exchange of mercury in a subtropical coniferous forest in south China. J Geophys Res Biogeosci 121(7):2006–2016
- Malinovsky D, Vanhaecke F (2011) Mercury isotope fractionation during abiotic transmethylation reactions. Int J Mass Spectrom 307:214–224
- Mélières M, Pourchet M, Dominique PC, Gaucher P (2003) Mercury in canopy leaves of French Guianain remote areas. Sci Total Environ Chem Geol 311:261–267
- Millhollen AG, Obrist D, Gustin MS (2006) Mercury accumulation in grass and forb species as a function of atmospheric carbon dioxide concentrations and mercury exposures in air and soil. Chemosphere 65:889–897
- Perrot V, Bridou R, Pedrero Z, Guyoneaud R, Monperrus M, Amouroux D (2015) Identical Hg isotope mass dependent fractionation signature during methylation by sulfate-reducing bacteria in sulfate and sulfate-free environment. Environ Sci Technol 49:1365–1373
- Poissant L, Pilote M, Yumvihoze E, Lean D (2008) Mercury concentrations and foliage/atmosphere fluxes in a maple forest

ecosystem in Quebec, Canada. J Geophys Res Atmos 113:D10307

- Rea AW, Lindberg SE, Scherbatskoy T, Keeler GJ (2002) Mercury accumulation in foliage over time in two northern mixedhardwood forests. Water Air Soil Pollut 133:49–67
- Richardson J, Friedland A (2015) Mercury in coniferous and deciduous upland forests in northern New England, USA: implications of climate change. Biogeosciences 12:6737–6749
- Selin NE (2009) Global Biogeochemical Cycling of Mercury: A Review. Annu Rev Environ Resour 34:43–63
- Selvendiran P, Driscoll CT, Montesdeoca MR, Bushey JT (2008) Inputs, storage, and transport of total and methyl mercury in two temperate forest wetlands. J Geophys Res 113:1–15
- Siwik EI, Campbell LM, Mierle G (2009) Fine-scale mercury trends in temperate deciduous tree leaves from Ontario, Canada. Sci Total Environ 407:6275–6279
- Song L, Zhang Y-J, Chen X, Li S, Lu H-Z, Wu C-S et al (2015) Water relations and gas exchange of fan bryophytes and their adaptations to microhabitats in an Asian subtropical montane cloud forest. J Plant Res 128:573–584
- Sun R, Enrico M, Heimburger LE, Scott C, Sonke JE (2013) A double-stage tube furnace-acid-trapping protocol for the preconcentration of mercury from solid samples for isotopic analysis. Anal Bioanal Chem 405:6771–6781
- Sun GY, Sommar J, Feng XB, Lin CJ, Ge MF, Wang WG et al (2016) Mass-dependent and -independent fractionation of mercury isotope during gas-phase oxidation of elemental mercury vapor by atomic Cl and Br. Environ Sci Technol 50:9232–9241
- Sun L, Lu B, Yuan D, Hao W, Zheng Y (2017) Variations in the isotopic composition of stable mercury isotopes in typical mangrove plants of the Jiulong estuary, SE, China. Environ Sci Pollut Res 24:1459–1468
- Tan Z-H, Zhang Y-P, Schaefer D, Yu G-R, Liang N, Song Q-H (2011) An old-growth subtropical Asian evergreen forest as a large carbon sink. Atmos Environ 45:15481554
- Wang D, Shi X, Wei S (2003) Accumulation and transformation of atmospheric mercury in soil. Sci Total Environ 304:209–214
- Wang C, Yan H, Liu H, Feng X, Wang J (2010) The method of rapidly measuring total mercury in solid samples using Lumex analytical equipment. Earth Environ (in chinese) 38:378–382 (in chinese)
- Wang X, Luo J, Yin R, Yuan W, Lin C-J, Sommar J et al (2017) Using mercury isotopes to understand mercury accumulation in the montane forest floor of the Eastern Tibetan Plateau. Environ Sci Technol 51:801–809
- Wang X, Yuan W, Lin C-J, Zhang L, Zhang H, Feng X (2019) Climate and vegetation as primary drivers for global mercury storage in surface soil. Environ Sci Technol 53:10665–10675
- Wiederhold JG, Cramer CJ, Daniel K, Infante I, Bourdon B, Kretzschmar R (2010) Equilibrium mercury isotope fractionation between dissolved Hg(II) species and thiol-bound Hg. Environ Sci Technol 44:4191–4197
- Xie S, Liu W, Li S, Guo P (1996) Preliminary study on the biomass of middle-mountain moist evergreen broadleaved forests in Ailao Mountain. Yunnan Acta Phytoecol Sin 20:167–176
- Yang L, Sturgeon R (2009) Isotopic fractionation of mercury induced by reduction and ethylation. Anal Bioanal Chem 393:377–385

- Yang Y, Yanai RD, Montesdeoca M, Driscoll CT (2017) Measuring mercury in wood: challenging but important. Int J Environ Anal Chem 13(4):e0196293
- Yang Y, Yanai RD, Driscoll CT, Montesdeoca M, Smith KT (2018) Concentrations and content of mercury in bark, wood, and leaves in hardwoods and conifers in four forested sites in the northeastern USA. PLoS ONE 13:e0196293
- Yin R, Feng X, Delphine F, Shi W, Zhao Z, Wanng J (2010) High precision determination of mercury isotope ratios using online mercury vapor generation system coupled with multicollector inductively coupled plasma-mass spectrometer. Chin J Anal Chem 38:929–934 (in chinese)
- Yin R, Feng X, Wang J, Bao Z, Yu B, Chen J (2013a) Mercury isotope variations between bioavailable mercury fractions and total mercury in mercury contaminated soil in Wanshan Mercury Mine SW, China. Chem Geol 336:80–86
- Yin RS, Feng XB, Meng B (2013b) Stable mercury isotope variation in rice plants (*Oryza sativa* L.) from the Wanshan Mercury Mining District, SW, China. Environ Sci Technol 47:2238–2245
- Yin R, Feng X, Chen J (2014) Mercury stable isotopic compositions in coals from major coal producing fields in China and their geochemical and environmental implications. Environ Sci Technol 48:5565–5574
- You CX (1983) Classification of vegetation in Xujiaba region in Ailao Mts. In: Wu ZY (ed) Research of forest ecosystem on Ailao Mountain, Yunnan. Yunnan Science and Technology Press, Kunming, pp 74–117
- Yu B, Fu X, Yin R, Zhang H, Wang X, Lin C-J et al (2016) Isotopic composition of atmospheric mercury in China: new evidence for sources and transformation processes in air and in vegetation. Environ Sci Technol 50:9262–9269
- Yuan W, Sommar J, Lin C-J, Wang X, Li K, Liu Y et al (2019) Stable isotope evidence shows re-emission of elemental mercury vapor occurring after reductive loss from Foliage. Environ Sci Technol 53:651–660
- Zheng W, Hintelmann H (2009) Mercury isotope fractionation during photoreduction in natural water is controlled by its Hg/DOC ratio. Geochim Cosmochim Acta 73:6704–6715
- Zheng W, Hintelmann H (2010a) Isotope fractionation of mercury during its photochemical reduction by low-molecular-weight organic compounds. J Phys Chem A 114:4246–4253
- Zheng W, Hintelmann H (2010b) Nuclear field shift effect in isotope fractionation of mercury during abiotic reduction in the absence of light. J Phys Chem A 114:4238–4245
- Zheng W, Obrist D, Weis D, Bergquist BA (2016) Mercury isotope compositions across North American forests Global Biogeochem. Cycles 30:1475–1492
- Zheng W, Demers JD, Lu X, Bergquist BA, Anbar AD, Blum JD et al (2019) Mercury stable isotope fractionation during abiotic dark oxidation in the presence of thiols and natural organic matter. Environ Sci Technol 53:1853–1862
- Zhou J, Wang Z, Sun T, Zhang H, Zhang X (2016) Mercury in terrestrial forested systems with highly elevated mercury deposition in southwestern China: the risk to insects and potential release from wildfires. Environ Pollut 212:188–196
- Zhou J, Wang Z, Zhang X, Gao Y (2017) Mercury concentrations and pools in four adjacent coniferous and deciduous upland forests in Beijin, China. J Geophys Res Biogeosci 122:1260–1274